

# Tubercular Meningitis Presenting as Multiple Cranial Nerve Palsy

RUPESH SHIVKUMAR PARATI<sup>1</sup>, PONVIJAYA MUTHUSWAMY YADAV<sup>2</sup>, VIJAYASHREE S GOKHALE<sup>3</sup>, KEYURI MEHTA<sup>4</sup>, CHIRAG GURBANI<sup>5</sup>



## ABSTRACT

Tubercular Meningitis (TBM) is caused due to the seeding of meninges with *Mycobacterium tuberculosis* and about one-third of the world's population is infected with it. The risk of contracting the infection increases in immunocompromised states which includes diabetes mellitus, chronic use of steroids, chronic alcoholism and Human Immunodeficiency Virus (HIV) co-infection. Here, a case of 47-year-old female has been presented, who reported to the Department of Medicine with complaints of weakness over the left side of the face associated with diminished hearing and tinnitus in the left ear and subjective vertigo since 20 days. Initially, all the above involvement was attributed to bell's palsy but no improvement was seen with its treatment. Incidentally, she was found to be HIV positive and lumbar puncture was done and eventually, she was diagnosed as a case of TBM. The patient was successfully treated with antitubercular medications and supportive measures and she responded well. She was discharged with antitubercular medications with regular follow-up in view of starting antiretroviral therapy after 6 weeks.

**Keywords:** Bell's palsy, Immunocompromised state, Tuberculosis

## CASE REPORT

A 47-year-old female patient reported to the Department of Medicine with complaints of weakness over the left side of the face from past 20 days associated with difficulty in chewing food and drooling of saliva from the left angle of the mouth. She also complained of difficulty in closing the left eye while sleeping. She had slurring of speech which was noticed by her son. There was also diminished hearing and tinnitus in the left ear along with subjective vertigo, which was intermittent since 20 days. She gave a history of weight loss of 10 kg in past 2 months. No significant past history was present. No history of similar complaints in the family members were present.


On examination, she was afebrile, pulse rate was 100 beats/minute, and blood pressure was 110/70 mmHg. On physical examination there was no pallor, icterus, cyanosis, clubbing, lymphadenopathy or oedema. Deviation of angle of the mouth towards right side was seen [Table/Fig-1].



[Table/Fig-1]: Deviation of angle of the mouth towards right side.

The visual acuity in the left eye was reduced to hand movement close to face. Cardiovascular, respiratory and per abdominal examination was within normal limits. Hence, with the above findings, it was concluded that the left 2<sup>nd</sup>, 7<sup>th</sup> and 8<sup>th</sup> cranial nerves were affected. Initially, the findings were attributed to bell's palsy and the patient was referred to Neurology and Ear, Nose and Throat Department

and she was started on tablet acyclovir 400 mg four times a day and injection dexamethasone 8 mg intravenous (i.v) three times a day with tapering dose. No improvement was seen after a week. On routine blood investigations, she was found to be HIV positive. The Cluster of Differentiation 4 (CD4) count was 98 cells/mm<sup>3</sup>. A review was done with a Neurologist and Magnetic Resonance Imaging (MRI) of the brain was done. The MRI brain was suggestive of Tubercular Meningitis (TBM) with multiple tubercular granulomatous lesions and involvement of proximal portion of 7<sup>th</sup> cranial nerve in the left Cerebellopontine angle [Table/Fig-2a,b,3].



[Table/Fig-2a,b]: Magnetic Resonance Imaging (MRI) brain (Fluid-Attenuated Inversion Recovery (FLAIR) sequence) shows a lesion in the left pons.

[Table/Fig-3]: MRI brain (T1 weighted sequence) shows multiple tubercular granulomatous lesions in the right parieto-occipital and left occipital region. (Images from left to right)

Lumbar puncture was done which showed lymphocytic predominant pleocytosis with elevated proteins and low glucose. Adenosine Deaminase (ADA) test was positive which was 24 U/L. Cartridge Based Nucleic Acid Amplification Test (CBNAAT) was negative. She was started on weight-based Antitubercular Therapy (AKT 4), tablet cotrimoxazole 800 mg OD, tablet aspirin 150 mg OD and steroids were continued. The patient improved within 10 days of starting the treatment and was discharged on AKT with regular follow-up in view of starting Antiretroviral Therapy (ART) after 6 weeks.

## DISCUSSION

Infection with *Mycobacterium tuberculosis* is a common complication of HIV infection. Individuals with HIV infection are more prone to develop pulmonary as well as extrapulmonary tuberculosis which can either be due to increased susceptibility of infection or

reactivation of latent tuberculosis [1]. The risk of extrapulmonary tuberculosis in a HIV infected individual increases with the declining CD4+ count usually less than 200/mm<sup>3</sup>. The most common cranial mononeuropathy in HIV patients involves the Facial nerve which usually occurs around the time of seroconversion. Facial paralysis in HIV infected patients does not clinically differ from typical bell's palsy, as seen in the present case. The neurological manifestations of Tuberculosis include TBM, tuberculous granulomas, or tuberculous brain abscess [2,3]. TBM contributes to high mortality rates in HIV patients, especially in developing countries, due to late diagnosis. Clinical manifestations are usually non specific and may mimic other opportunistic infections of Central Nervous System (CNS) such as cryptococcosis, neurosyphilis, and bacterial meningitis [4].

Berenguer J et al., reported that the frequent presenting symptoms of TBM in a HIV infected individual were fever, headache and altered consciousness [5]. However, no such symptoms were present in this case. Another study conducted by Katrak SM et al., reported that most of the HIV positive patients with TBM had cognitive impairment on presentation [6]. However, no cognitive impairment was observed in this case. A similar case reported by Kama S et al., reported a case of HIV positive patient with multiple cranial nerve palsies including sixth, ninth, tenth and twelfth nerve [7].

In a study conducted by Dian S et al., various findings of TBM in brain imaging was meningeal enhancement, tuberculomas, brain infarction and hydrocephalus [8]. The present case had all findings except hydrocephalus. Therefore, a high index of suspicion should always be considered, and lumbar puncture along with radiological interventions should be considered in patients with HIV infection.

Delayed initiation of ART is associated with increased risk of mortality hence tuberculosis should always be considered as a differential diagnosis in patients with HIV. It has also been reported that early initiation of antiretroviral therapy among People Living with HIV (PLHIV) with TB meningitis is associated with adverse effects, hence it is safer to delay the initiation of ART [9]. TB Immune Reconstitution Inflammatory Syndrome (IRIS) manifests as paradoxical worsening of existing TB infection or development of new infection after initiation of antiretroviral therapy in a patient with tuberculosis [10].

HIV infected individuals who present with features of bell's palsy should always undergo a lumbar puncture to rule out tubercular infection. The goal is early diagnosis and initiation of antitubercular treatment in these patients.

## CONCLUSION(S)

Tubercular infection was suspected in the patient as no improvement was seen with treatment of bell's palsy. Tuberculosis complicates the disease course of HIV infection. Hence, early detection and initiation of treatment with ART is necessary to reduce the incidence of HIV related mortality.

## Acknowledgement

Authors are thankful to the guide Dr. Vijayashree S. Gokhale, Professor, Department of Medicine, Dr. D.Y. Patil Medical College for her support and guidance.

## REFERENCES

- [1] Bruchfeld J, Correia-Neves M, Källenius G. Tuberculosis and HIV coinfection. *Cold Spring Harb Perspect Med.* 2015;5(7):a017871.
- [2] De Cock KM, Soro B, Coulibaly IM, Lucas SB. Tuberculosis and HIV infection in sub-Saharan Africa. *JAMA.* 1992;268(12):1581-87.
- [3] Pujari S. Opportunistic infections in HIV, *API Textbook of Medicine*, 11<sup>th</sup> edition, page 377-382.
- [4] Croda MG, Vidal JE, Hernández AV, Dal Molin T, Gualberto FA, de Oliveira AC. Tuberculous meningitis in HIV-infected patients in Brazil: clinical and laboratory characteristics and factors associated with mortality. *International Journal of Infectious Diseases.* 2010;14(7)
- [5] Berenguer J, Moreno S, Laguna F, Vicente T, Agrados M, Ortega A, et al. Tuberculous meningitis in patients infected with the human immunodeficiency virus. *N Engl J Med.* 1992;326(10):668-72.
- [6] Katrak SM, Shembalkar PK, Bijwe SR, Bhandarkar LD. The clinical, radiological and pathological profile of tuberculous meningitis in patients with and without human immunodeficiency virus infection. *J Neurol Sci.* 2000;181(1-2):118-26.
- [7] Kama S, Biswas J, Kumarasamy N, Sharma P. Multiple cranial nerve palsy in an HIV-positive patient. *Indian J Ophthalmol.* 2001;49:118-20.
- [8] Dian S, Hermawan R, van Laarhoven A, Immaculata S, Achmad TH, Ruslami R, et al. Brain MRI findings in relation to clinical characteristics and outcome of tuberculous meningitis. *PLoS One.* 2020;15(11)
- [9] Török ME, Yen NT, Chau TT, Mai NT, Phu NH, Mai PP, et al. Timing of initiation of antiretroviral therapy in human immunodeficiency virus (HIV)-associated tuberculous meningitis. *Clin Infect Dis.* 2011;52(11):1374-83.
- [10] Mishra V, Harbada R, Sharma A, Narute S. Tuberculosis-immune reconstitution inflammatory syndrome. *Journal of HIV and Human Reproduction.* 2014;2(2):63.

### PARTICULARS OF CONTRIBUTORS:

1. Resident, Department of Internal Medicine, Dr. D.Y. Patil Medical College, Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.
2. Resident, Department of Internal Medicine, Dr. D.Y. Patil Medical College, Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.
3. Professor, Department of Internal Medicine, Dr. D.Y. Patil Medical College, Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.
4. Resident, Department of Internal Medicine, Dr. D.Y. Patil Medical College, Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.
5. Resident, Department of Internal Medicine, Dr. D.Y. Patil Medical College, Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.

### NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Ponvijaya Muthuswamy Yadav,  
Resident, Department of Internal Medicine, Dr. D.Y. Patil Medical College,  
Dr. D.Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India.  
E-mail: ponvijayamuthuswamy@gmail.com

### PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Jan 20, 2022
- Manual Googling: Jan 27, 2022
- iThenticate Software: Feb 14, 2022 (14%)

### ETYMOLOGY: Author Origin

### AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: **Jan 19, 2022**

Date of Peer Review: **Feb 04, 2022**

Date of Acceptance: **Mar 29, 2022**

Date of Publishing: **Apr 01, 2022**