


Association of Human Leukocyte Antigen Gene Variants rs13192471 and rs6457617 with Rheumatoid Arthritis Susceptibility: A Case-control Study from North-western India

INDERMOHAN SINGH SANDHU¹, HEMENDER SINGH², ANUPAMA MAHAJAN³, GURINDER MOHAN⁴, SWARKAR SHARMA⁵, PRAKASH SINGH⁶, SIMRANPREET KAUR⁷, AMARJIT SINGH BHANWER⁸



ABSTRACT

Introduction: Rheumatoid Arthritis (RA) is a systemic, chronic, inflammatory, and autoimmune disease which is characterised by the progressive ruination of joint structures. The Human Leukocyte Antigens (HLA) genes: HLA-DRB1 and HLA-DQB1, belonging to HLA family, presented on human chromosome six, are involved in the immune system. Various studies involving Genome-wide Association Studies (GWAS), meta-analysis, and replication studies have shown the association of HLA-DRB1 variant rs13192471 and HLA-DQB1 variant rs6457617 with RA susceptibility in different population groups.

Aim: To perform a replication case-control based association study of variants rs13192471 and rs6457617, in order to determine their association with RA, in an independent cohort from the population of North-western India.

Materials and Methods: The case-control study conducted at Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India, duration of the study was,

from May 2019 to June 2022. In this study, Deoxyribonucleic Acid (DNA) was isolated from 188 RA cases and 310 healthy controls, followed by qualitative and quantitative analyses. The genotyping was performed on a Real Time-polymerase Chain Reaction (RT-PCR) using TaqMan Allele Discrimination Assay. Statistical power of the study was estimated using PS: Power and sample size calculator version 3.1.

Results: The statistical analysis of the genotyping data showed a significant association ($p=0.005$) of variant rs13192471 with RA susceptibility, whereas the variant rs6457617 did not show an association ($p=0.275$) with RA in the studied population cohort.

Conclusion: The present study successfully replicated the association of HLA-DRB1 variant rs13192471 in the population group of North-western India. It is pertinent to screen these variants for RA susceptibility in other population groups of India before their use as potential genetic biomarkers in the Indian population.

Keywords: Autoimmune diseases, Genome wide association study, Genotyping, TaqMan assay

INTRODUCTION

The rheumatoid arthritis is a chronic systemic autoimmune disease with immunologically arbitrated inflammation of synovia-lined joints that can significantly disrupt joint construction and task [1]. RA is an exemplar of an intricate genetic state with undetermined genetic diversity and origin, and it almost certainly involves several genes [2]. A persistent inflammatory joint illness RA is known to be caused by reactions of autoimmunity to Anti-citrullinated Peptide Antibody (ACPA) and Immunoglobulin G (IgG), frequently known to be Rheumatoid Factor (RF) [3]. Besides RF, RA patients often exhibit antibodies peculiar to other proteins, such as type II collagen (COL-II) and Human Cartilage glycoprotein 39 (HC gp-39) [4]. Regular demonstration of RA is generally polyarthritis with firmness, ache, and protuberance of multiple joints in a symmetric pattern [5].

The human Major Histocompatibility Complex (MHC) system genes, also known as the HLA family, are one of the most polymorphic genetic systems and control effector immunity units [6]. The heritability of RA is approximately 60%, to which HLA contributes around 11-37% [7]. *HLA-DRB1* and *HLA-DQB1* genes are located on chromosome 6, play an essential role in the immune system, and are well known for playing a role in RA pathogenesis [8]. A strong connection between RA and HLA alleles has long been reported.

The DR and DQ loci of the *HLA* gene have been reported in strong linkage disequilibrium and associated with RA in many studies of Caucasian and East asian populations [9]. GWAS of RA also exhibited remarkable associations with various Single Nucleotide Polymorphisms (SNPs) near the *HLA-DRB1* gene with RA in European, Korean and Japanese populations. The significant SNPs with this association are rs7765379, rs6457617, rs660895 and rs13192471 [10-12]. HLA variant rs6457617 was also associated with RA in Spanish and Han Chinese populations [13,14]. HLA SNP rs6457617 was associated with RA susceptibility, while SNP rs13192471 showed no association with RA in Tunisian [12]. Indian populations are highly diverse. A study from North-eastern India showed an association of SNP rs13192471 in susceptibility and severity of RA, whereas SNP rs6457617 did not [15]. However, another study showed that both HLA SNPs, rs13192471 and rs6457617, are associated with a significant risk for RA in the Northern Indian population [16] which indicate genetic heterogeneity in RA.

As there is significant genetic heterogeneity in RA among different world populations, replication analysis studies to find the association of various susceptibility loci with RA in multiple populations is essential, making screening in the North-western Indian population critical. With this objective, the present study was performed

in a case-control association designed to screen HLA variants rs13192471 and rs6457617 in the North-western Indian population cohort.

MATERIALS AND METHODS

The present case-control study was conducted at Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar in association with Sri Mata Vaishno Devi University, Katra, Jammu and Kashmir, duration of the study was, from May 2019 to June 2022. The study was initiated after the due approval by the Institutional Ethics Committee (Ref no. Patho 339/19, dated 06/04/2019). After getting the written informed consent of the participants, a venous blood sample of 2 mL was obtained and transferred to Ethylenediaminetetraacetic Acid (EDTA) vials to avoid coagulation. The samples were transported to a laboratory, in an icebox and kept at -20°C until the Deoxyribonucleic Acid (DNA) had been extracted. Participants included 188 ACR/EULAR (American College of Rheumatology/European League Against Rheumatism) Criterion confirmed RA cases, and 310 age and sex matched healthy controls.

Inclusion criteria: The minimum age for cases to include in this study was 18 years to avoid inclusion of juvenile RA cases in this study with no upper limit of age for the inclusion of cases. Same age criteria were kept for controls with disease free status related to any autoimmune disease, and any disease related to bone, articular cartilage, or endocrine system.

Exclusion criteria: For both cases and controls, individuals with conditions like cancer, diabetes mellitus, neurological disorders, thyroid dysfunction, infectious disease, arthritis other than RA for cases and females with pregnancy and nursing mothers were excluded.

Study Procedure

The isolation of DNA is done using a MagGenome® XpressDNA blood kit. Then agarose gel electrophoresis was performed to determine the quality of isolated DNA samples. The concentration of the DNA samples was determined using NanoDrop™ 2000/2000c (Thermo Scientific). The 50 µL dilutions (10 ng/µL) of DNA samples were prepared by adding nuclease-free water.

Genotyping of variant rs13192471 and rs6457617: Using the TaqMan allele discrimination test, the SNP genotyping of variants was assessed in the studied population group. Taqman® SNP genotyping assay from Applied Biosystems (Thermo Fischer Scientific, Pleasanton, CA, USA) was used for genotyping the variants rs13192471 and rs6457617 using Mx3005p Agilent RT-PCR (Stratagene Agilent Technologies, Waldbronn, Germany). The Taqman genotyping assay was diluted using TE (Tris EDTA) from 40X to 20X as directed by the manufacturer. The genotyping was performed using a 96-well plate with 93 samples and three Non Template Controls (NTC). The overall amount of the PCR reaction mix was 10 µL, including 3 µL of DNA concentration of 10 ng/µL. Rest of the constituents included 1 µL 1X PCR buffer, 0.25 µL 0.5X of TaqMan assay, 0.2 µL 0.2 mM of dNTPs, 0.1 µL 0.3 U of Taq Polymerase, 5.45 µL of distilled water was used to make up the volume of the reaction. The cycling conditions consist of a 4-minute hold at 95°C, 40 cycles at 15-second intervals at 95°C, and one minute at 60°C with endpoint detection. The RT-PCR post-PCR detection equipment was used to assess the allele-specific fluorescence. Reruns were performed to determine final inclusion for samples that failed genotyping or had ambiguous genotypes.

STATISTICAL ANALYSIS

Statistical power of the study was estimated using PS: Power and sample size calculator version 3.1. The analysis of genotyping data was performed in PLINK version 1.09. An Odds Ratio (OR)

was calculated to determine the chance of an outcome in response to exposure (allele or genotype). The value and 95% Confidence Interval (CI), which gives the relationship or assesses the likelihood of an outcome, were computed together. The OR indicates how frequently the patients are more at risk than controls. The Hardy Weinberg Equilibrium (HWE) was also calculated using the Chi-square test.

RESULTS

Most of the RA cases in this study cohort were females (146/188, 77.65%). For enrolled subjects, the average age was 48.43±12.08 years. The average age of onset of disease was 44.49±10.96 years. Of 188 RA cases, 68.61% of the patients were seropositive for RF. This study group's average Body Mass Index (BMI) was 24.72±3.78. The middle Waist-Hip Ratio (WHR) was 0.88±0.072. Only 3.19% of patients were smokers, and 5.85% took alcohol regularly. Around 62% of the patients were taking a purely vegetarian diet.

SNP Analyses

Genotypic analyses of single nucleotide polymorphism (SNP)

rs6457617: The allelic distribution of variant rs13192471 was evaluated. The C allele was observed as a minor allele with a frequency of 0.18 (18%) in cases and 0.11 (11%) in controls in the studied cohort. The variant rs13192471 showed a significant association (p-value 0.005) with RA susceptibility and the T allele as a risk allele for RA with the OR of 1.77, 95% CI (1.18-2.64), in the present studied cohort. The allelic distribution of variant rs13192471 of *HLADRB1* was in not in accordance with HWE in both the cases as well as controls.

In order to further evaluate and increase the power of the study and overcome confounding factors effects, logistic regression analyses were adopted with the recessive (CT+TT vs CC) and dominant model (CT+CC vs TT), where TT is wild homozygous genotype, CT is heterozygous genotype and CC is mutant homozygous genotype. In the dominant model, the OR obtained was 1.96 (1.23-3.14 at 95% CI), p=0.004, adjusted with age, gender, BMI and ethnicity. The results showed a significant association of variant rs13192471 with RA in the population belonging to Northern India. The frequency of TT, CT and CC genotype was found to be 69.18% vs 81.51%, 25.34% vs 14.72% and 5.48% vs 3.77% in the RA cases and controls respectively. A Chi-square test of independence was performed to examine the relation between three genotypes between cases and controls. The result came to be significant with the p-value 0.016 [Table/Fig-1].

rs13192471	Frequency in cases (%)	Frequency in controls (%)	OR (95% CI)	p-value
Genotype				
Homozygous wild TT	69.18	81.51	1.77 (1.18-2.64)	0.016
Heterozygous variant CT	25.34	14.72		
Homozygous variant CC	5.48	3.77		
Allele frequency in decimal				
T allele	0.82	0.89	1.77 (1.18-2.64)	0.005
C allele	0.18	0.11		
Genetic model				
Dominant model (CT+CC vs TT)			1.96 (1.23-3.14)	0.004
Recessive model (CT+TT vs CC)			1.48 (0.57-3.83)	0.418

[Table/Fig-1]: Genotype and allele frequencies of rs13192471 polymorphism in RA patients and controls.

CI: Confidence level; OR: Odds ratio; p-value <0.05 is considered as significant

Genotypic analyses of single nucleotide polymorphism

rs6457617: The allelic distribution of variant rs6457617 was

evaluated. The T allele was observed as a minor allele with a frequency of 0.38 (38%) in cases and 0.34 (34%) in controls in the studied cohort. The variant rs6457617 showed a non significant association (p-value 0.275) with RA susceptibility with an OR of 1.18, 95% CI (0.87-1.60), in the present studied cohort. The allelic distribution of variant rs6457617 of *HLADQB1* was in not in accordance with HWE in both the cases as well as controls. In order to further evaluate and increase the power of the study and overcome confounding factors effects, logistic regression analyses were adopted with dominant model (CT+TT vs CC) and recessive models (CT+CC vs TT), where CC is homozygous wild genotype, CT is heterozygous genotype and TT is homozygous mutant genotype. The results showed no significant association of variant rs6457617 with RA in the population belonging to Northern India (p>0.05). The frequency of CC, CT and TT genotype was found to be 44.12% vs 48%, 36.03% vs 36% and 19.85% vs 16% in the RA cases and controls respectively. A Chi-square test of independence was performed to examine the relation between these three genotypes in cases and controls. The result came to be non significant with the p-value 0.584 [Table/Fig-2].

rs6457617	Frequency in cases (%)	Frequency in controls (%)	OR (95% CI)	p-value
Genotype				
Homozygous wild CC	44.12	48	1.18 (0.87-1.60)	0.584
Heterozygous variant CT	36.03	36		
Homozygous variant TT	19.85	16		
Allele frequency in decimal				
C allele	0.62	0.66	1.18 (0.87-1.60)	0.275
T allele	0.38	0.34		
Genetic model				
Dominant model (CT+TT vs CC)			1.17 (0.77-1.77)	0.457
Recessive model (CT+CC vs TT)			1.30 (0.77-2.21)	0.330

[Table/Fig-2]: Genotype and allele frequencies of rs6457617 polymorphism in RA patients and controls.
CI: Confidence level; OR: Odds ratio; p-value <0.05 is considered as significant

DISCUSSION

The RA is a long-term autoimmune disorder with a world prevalence of approximately 0.46% [17]. India's majority distribution is found to vary among various geographical regions and ranges from 0.28% to 0.75% [18]. Various GWAS and meta-analyses in Asian populations came out with results which link multiple genes/loci of known immune function with RA susceptibility [19]. The disease has been explored for its genetic susceptibility to various genes and its SNPs across diverse populations using multiple study designs. Findings from these studies signify genetic heterogeneity in RA susceptibility across different populations [16]. The involvement of genes from the HLA region in RA susceptibility and severity has been well-known [9], especially the HLA class II genes which include *DQA1*, *DQB1* and *DRB1* [20]. The SNPs from the HLA region, rs13192471 and rs6457617, were most widely reported across various populations across the world with heterogeneity. Thus, the variants were chosen to study the association with RA in a population cohort from North-western India.

Interestingly, the variant rs13192471 was found to be significantly associated with risk (p-value=0.005) in present study population. At the same time, the variant rs6457617 was not significantly associated with RA susceptibility (p-value=0.275) in this population cohort. Our findings showed similar results as observed in the North-eastern Indian population [15] and partial overlap with the study from populations corresponding to Northern India [16] and the European population [10-12] but contrasting to results from Tunisian, Spanish and Han Chinese populations, where they

established a significant association of rs6457617 [12-14]. A peculiar finding of deviation in the genotype frequencies for both the variants from the HWE was observed in the study. This could be attributed to the polymorphic nature of the HLA locus and another possibility that the region might be under natural selection and with bias in allele frequencies of the variants, thereby not following HWE.

Limitation(s)

The study group comprised of participants from North-western India only. As Indian population is very diverse, replication and exploration in other population cohorts is critically needed.

CONCLUSION(S)

This is the first replication study evaluating the association of *HLA-DRB1* variant rs13192471 and *HLA-DQB1* variant rs6457617 in the population group of North-western India. The present study highlights the need to replicate the study with increased sample size and investigate other genes in the HLA region with RA susceptibility. Moreover, with high genetic diversity and thus anticipated genetic heterogeneity, such replication studies are warranted in different population groups of India. They will strengthen the identification of potential genetic biomarkers for the various Indian populations and contribute towards futuristic goals of polygenic risks and personalised medicine.

REFERENCES

- [1] Kay J, Upchurch KS. ACR/EULAR 2010 rheumatoid arthritis classification criteria. *Rheumatology (Oxford)*. 2012;51(Suppl 6):vi5-9.
- [2] Weyand CM, Goronzy JJ. Association of MHC and rheumatoid arthritis: HLA polymorphisms in phenotypic variants of rheumatoid arthritis. *Arthritis Research & Therapy*. 2000;2(3):01-05.
- [3] Abbasi M, Mousavi MJ, Jamalzei S, Alimohammadi R, Bezvan MH, Mohammadi H, et al. Strategies toward rheumatoid arthritis therapy; the old and the new. *Journal of Cellular Physiology*. 2019;234(7):10018-31.
- [4] Nepomnyashchikh T, Antonets D, Shchelkunov S. Gene therapy of arthritis. *Russian Journal of Genetics*. 2016;52(6):543-56.
- [5] Harris ED Jr. Rheumatoid arthritis. Pathophysiology and implications for therapy. *The New England Journal of Medicine*. 1990;322(18):1277-89.
- [6] Golovanova O, Kononov V, Shevchenko A, Smolnikova M. Frequencies of the DRB1, DQA1, DQB1 and TNFA alleles in immigrant population of west Siberia. *Russian Journal of Genetics*. 2009;45(8):981-86.
- [7] Kurko J, Besenyei T, Laki J, Glant TT, Mikecz K, Szekanecz Z. Genetics of rheumatoid arthritis-a comprehensive review. *Clin Rev Allergy Immunol*. 2013;45(2):170-79.
- [8] Wysocki T, Olesinska M, Paradowska-Gorycka A. Current understanding of an emerging role of HLA-DRB1 gene in rheumatoid arthritis-from research to clinical practice. *Cells*. 2020;9(5):1127.
- [9] Van Dongen V, Holoshitz J. Human leukocyte antigen-disease associations in rheumatoid arthritis. *Rheumatic Disease Clinics*. 2017;43(3):363-76.
- [10] Wellcome Trust Case Control C. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature*. 2007;447(7145):661-78.
- [11] Plenge RM, Seielstad M, Padyukov L, Lee AT, Remmers EF, Ding B, et al. TRAF1-C5 as a risk locus for rheumatoid arthritis-a genomewide study. *The New England Journal of Medicine*. 2007;357(12):1199-209.
- [12] Achour Y, Ben Hamad M, Chaabane S, Rebai A, Marzouk S, Mahfoudh N, et al. Analysis of two susceptibility SNPs in HLA region and evidence of interaction between rs6457617 in HLA-DQB1 and HLA-DRB1*04 locus on Tunisian rheumatoid arthritis. *Journal of Genetics*. 2017;96(6):911-18.
- [13] Julia A, Ballina J, Canete JD, Balsa A, Tornero-Molina J, Naranjo A, et al. Genome-wide association study of rheumatoid arthritis in the Spanish population: KLF12 as a risk locus for rheumatoid arthritis susceptibility. *Arthritis and Rheumatism*. 2008;58(8):2275-86.
- [14] Li H, Hu Y, Zhang T, Liu Y, Wang Y, Yang T, et al. Replication of british rheumatoid arthritis susceptibility Loci in two unrelated chinese population groups. *Clinical & Developmental Immunology*. 2013;2013:891306.
- [15] Das S, Baruah C, Saikia AK, Bose S. Associative role of HLA-DRB1 SNP genotypes as risk factors for susceptibility and severity of rheumatoid arthritis: A North-east Indian population-based study. *Int J Immunogenet*. 2018;45(1):01-07.
- [16] Prasad P, Kumar A, Gupta R, Juyal RC, Thelma BK. Caucasian and Asian specific rheumatoid arthritis risk loci reveal limited replication and apparent allelic heterogeneity in north Indians. *PLoS One*. 2012;7(2):e31584.
- [17] Almutairi K, Nossent J, Preen D, Keen H, Inderjeeth C. The global prevalence of rheumatoid arthritis: A meta-analysis based on a systematic review. *Rheumatol Int*. 2021;41(5):863-77.
- [18] Handa R, Rao UR, Lewis JF, Rambhad G, Shiff S, Ghia CJ. Literature review of rheumatoid arthritis in India. *International Journal of Rheumatic Diseases*. 2016;19(5):440-51.

- [19] Hollis-Moffatt JE, Chen-Xu M, Topless R, Dalbeth N, Gow PJ, Harrison AA, et al. Only one independent genetic association with rheumatoid arthritis within the KIAA1109-TENR-IL2-IL21 locus in Caucasian sample sets: Confirmation of association of rs6822844 with rheumatoid arthritis at a genome-wide level of significance. *Arthritis Res Ther.* 2010;12(3):R116.
- [20] Ortiz-Fernandez L, Martin J, Alarcon-Riquelme ME. A summary on the genetics of systemic lupus erythematosus, rheumatoid arthritis, systemic sclerosis, and sjogren's syndrome. *Clin Rev Allergy Immunol.* 2022.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Genetics, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India.
2. Research Associate, Department of Biotechnology, Sri Mata Vaishno Devi University, Katra, Jammu and Kashmir, India.
3. Professor, Department of Anatomy, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India.
4. Professor and Head, Department of Medicine, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India.
5. Assistant Professor, Department of Biotechnology, Sri Mata Vaishno Devi University, Katra, Jammu and Kashmir, India.
6. Senior Orthopaedics Consultant, Department of Orthopaedics, Parkash Hospital, Amritsar, Punjab, India.
7. Assistant Professor, Department of Genetics, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India.
8. Professor and Head, Department of Genetics, Sri Guru Ram Das Institute of Medical Sciences and Research, Amritsar, Punjab, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Indermohan Singh Sandhu,
Assistant Professor, Department of Genetics, SGRD Institute of Medical Sciences and Research, Mehta Road, Amritsar-143501, Punjab, India.
E-mail: imssandhu.gen@sgrdimr.in

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Feb 18, 2023
- Manual Googling: Mar 02, 2023
- iThenticate Software: Mar 25, 2023 (5%)

ETYMOLOGY: Author Origin**AUTHOR DECLARATION:**

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. No

Date of Submission: **Feb 14, 2023**Date of Peer Review: **Mar 11, 2023**Date of Acceptance: **Apr 05, 2023**Date of Publishing: **May 01, 2023**