

Immediate Effect of Kinetic Chain Activation and Mulligan's Bent Leg Raise Technique in Patients with Asymptomatic Hamstring Tightness: A Randomised Clinical Trial

SANTOSH CHANNABASAPA METGUD¹, SANTOSH D PATIL², OMKAR ANAND GAONKAR³

ABSTRACT

Introduction: Hamstring shortness is an important mechanical marker for hamstring strain injuries, often related to a muscle imbalance between the quadriceps and hamstrings. Muscle tightness is considered a limiting factor for optimal performance, including individuals' daily activities.

Aim: To compare the immediate effects of the Kinetic Chain Activation Technique (KCAT) and Mulligan's Bent Leg Raise (BLR) on hamstring tightness using the active knee extension test, diagnostic ultrasonography and the Sit and Reach Test (SRT) test.

Materials and Methods: A randomised clinical trial was conducted in the Orthopaedic Manual Therapy Department of KAHER Institute of Physiotherapy, Belagavi, Karnataka, India, from December 2022 to June 2023. In this trial, subjects with asymptomatic hamstring tightness were enrolled and randomised into two interventional groups: KCAT (n=15) and BLR (n=15). Kinetic chain activation was applied to Interventional

group A, while Mulligan's BLR was applied to Interventional group B. Measurements were taken of the hamstring muscle's length using the SRT test, the active knee extension test and diagnostic ultrasound to determine the immediate effects. To compare the length and thickness of the hamstring, AKT and SRT test within groups, the paired t-test was used. For between-group analysis, the independent t-test was applied.

Results: The mean age of the subjects was 23.8±0.47 years. There were 10 males and five females in group A and eight males and seven females in group B. Significant differences were identified in the thickness of the hamstring muscles in both groups when analysed within groups (p-value=0.001). However, there was no significant difference in the between-group analysis (p>0.001).

Conclusion: The study concluded that both Mulligan's BLR and KCAT were equally effective and could be used as adjuncts in improving hamstring flexibility.

Keywords: Active knee extension test, Fascia, Hamstring muscle, Sit and reach test

INTRODUCTION

Since the hamstring muscles are the primary muscles responsible for hip extension and knee flexion, they are essential for the regular performance of functional activities [1,2]. As a monoarticular muscle that shares a tendon with the biceps femoris, the hamstring is typically tested in conjunction with other hamstrings [3]. Hamstring shortening is a crucial mechanical marker for hamstring strain injuries and is often associated with an imbalance in the strength and stiffness of the hamstring and quadriceps muscles [4]. Muscle tension is believed to be a barrier to an individual's maximum performance, including daily activities [5].

Adaptive shortening of the muscle's non contractile and contractile components is referred to as "muscle tightness." Even under normal conditions, the bi-articular postural muscle known as the hamstring tends to shorten [6]. It is a superficial two-joint muscle that can become exceedingly tight, leading to a range of postural problems and muscular imbalances [7]. Clinical data indicate that hamstring tightness may be associated with alterations in the transverse spine curvatures during trunk flexion, affecting the lumbar pelvic rhythm [8]. Short hamstrings may increase the risk of spinal damage from mechanical stresses [9].

Tension in the fascia ultimately impacts the body's overall range of motion [10-15]. De-tension must be established in the fascia to relieve symptoms. De-tensioning the hamstring using K-CAT results in a decrease in hamstring tightness [16]. The K-CAT involves the activation and mobilisation of fascia along the fascial chains. The

idea behind a kinetic chain is that these joints and segments interact with one another during movement. When one joint moves, it sets off a series of events that affect the movement of nearby joints and segments. An integrated biomechanical task is the outcome of sequenced physiological muscle activations in the upper and lower extremities. A Mulligan stretching method called the BLR is used to treat tight hamstrings. It is a non invasive, helpful technique recommended when hamstring stiffness prevents the performance of the Straight Leg Raise (SLR) [17-19].

The current study aimed to fill the information gap by comparing the effectiveness of KCAT and BLR in terms of objective outcome measures that reflect changes in muscle and tissue levels.

MATERIALS AND METHODS

A randomised clinical trial was conducted in the Orthopaedic Manual Therapy Department of KAHER Institute of Physiotherapy, Belagavi, Karnataka, India, from December 2022 to June 2023. Institutional Ethical Committee approval was obtained (Ref No: KIPT/706/12/05/22) and the trial was registered under the Clinical Trial Registry - India (CTRI/2023/08/057073).


Inclusion criteria: Participants aged 18-25 years with asymptomatic hamstring tightness, regardless of gender, were included. Individuals with a lack of active knee extension greater than 15 degrees were also included [8].

Exclusion criteria: The exclusion criteria included: 1) Participants with recent musculoskeletal injuries; 2) Recent injuries around the hip and knee joints.

Sample size: The minimum sample size required for the execution of the research study was 30, with 15 participants in each group.

Study Procedure

Demographic data was obtained and all participants received the therapy. Participants were divided into two groups. Simple randomisation was performed using the sealed envelope method. Participants were blinded to the study and randomised into two groups: the KCAT group (group A) and the BLR group (group B). The Consolidated Standards of Reporting Trials (CONSORT) flow diagram for recruitment of participants has been given in [Table/Fig-1].



Intervention details:

Group A: (Hamstring Fascia Activation through KCAT Technique) [Table/Fig-2a]. The participant was positioned prone, lying on a plinth with the hip and knee in a neutral position. The therapist stood beside the leg to be tested. The therapist activated the hamstring fascia by tapping on the hamstring area for 7-8 seconds. After this, the patient was asked to flex the knee while the therapist applied pressure on the tendo Achillis until the patient could flex fully and then extend. This was done for 10 minutes. The same procedure was performed on the opposite lower extremity [20]. The duration of the intervention was two weeks, with three sessions per week, each lasting 15 minutes.

Group B: Mulligan's Bent Leg Raise (BLR) [Table/Fig-2b].



[Table/Fig-2]: a) Kinetic Chain Activation Technique (KCAT). b) Mulligan's Bent Leg Raise (BLR).

The therapist stood at the side of the subject. The participant's hip and knee were in 90° flexion, with the heel off the plinth. The therapist's position was a walking stance, lateral to the affected side. The participant held the plinth with the unaffected side and placed the hand on the affected side under their head and neck. The therapist's inner hand was placed underneath, proximal to the popliteal fossa. The therapist grasped the lower end of the thigh with both hands. Longitudinal traction was applied along the long axis of the femur, taking the hip into flexion until the first resistance was felt. If the subject complained of stretch pain, then a contract-relax technique was applied by asking the subject to gently push against the therapist's shoulder for five seconds. The leg could then be taken into a new pain-free range. If the subject experienced pain, the hip was moved into abduction or external rotation, or more traction was given before further hip flexion was added. The end position was maintained for about 20 seconds. This technique was repeated three times [21,22]. The duration of the intervention was two weeks, with three sessions per week, each lasting 15 minutes.

Preassessment of the thickness and length of the hamstring muscles was performed using ultrasonography, while the active knee extension test and SRT test were also applied. Post-assessment data were calculated immediately after the intervention.

Active knee extension test: The therapist was positioned to the side of the leg being evaluated. To minimise pelvic rotation, the opposite extremity was secured with a strap. The patient lay supine with their hips flexed at 90 degrees. The participant was instructed to extend their knee as far as possible and a universal goniometer was used to assess the result [23].

Sit and reach test: In addition to the subject or participant, the therapist occupied the following position: the patient sat on the floor with their dominant leg extended and the other leg flexed. Their bare feet were positioned flat against a box at a 90-degree angle to the floor. While reaching forward, the subjects performed the sit and reach with their palms toward the floor. If the participant experienced any pain or stiffness in their posterior thigh, they were instructed to stop. The distance travelled was measured in centimetres for the five seconds that this posture was held. The highest score from the two assessments taken before and after the intervention was used to analyse the final results [24].

STATISTICAL ANALYSIS

The collected data were summarised using descriptive statistics: frequency, percentage, mean and Standard Deviation (SD). To compare the length and thickness of the hamstring, AKT and SRT test within groups, the paired t-test was used. For comparisons between groups, the independent t-test was employed. The Shapiro-Wilk test was conducted to assess the normality of the data. A p-value of less than 0.05 was considered significant. Data were analysed using Statistical Package for the Social Sciences (SPSS) software (SPSS Inc.; Chicago, IL) version 29.0.10.

RESULTS

In the current study, 16 males and 14 females were included, with a mean age of 23.8±0.47 years and a mean Body Mass Index (BMI) of 24.2±0.56 kg/m². There was a significant difference between the pre- and post-treatment values in both groups [Table/Fig-3]. There

Groups	Times	Mean±SD	SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	16.19±1.17	1.17	1.51	0.46	3.26	13.025	0.001*
	Post	17.69±1.15	1.15					
Group-B	Pre	16.13±0.99	0.99	1.45	0.57	2.54	9.844	0.001*
	Post	17.59±0.91	0.91					

[Table/Fig-3]: Comparison of pretest and post-test scores of thickness of hamstring muscle in right leg using diagnostic ultrasonography within the two groups. Paired sample t-test

was a significant difference between the pre- and post-treatment values in both groups [Table/Fig-4].

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	16.48±1.20	1.30	0.65	1.99	7.961	0.001*
	Post	17.78±1.15					
Group-B	Pre	16.16±1.17	1.67	0.89	1.88	7.275	0.001*
	Post	17.83±0.87					

[Table/Fig-4]: Comparison of pretest and post-test scores of thickness of hamstring muscle in left leg using diagnostic ultrasonography within the two groups. Paired sample t-test

There was a significant difference between the pre- and post-treatment values in both groups as determined by the paired sample t-test [Table/Fig-5]. There was a significant difference between the pre- and post-treatment values in both groups as determined by the paired sample t-test [Table/Fig-6]. There was a significant difference between the pre- and post-treatment values in both groups as determined by the paired sample t-test [Table/Fig-7].

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	37.44±6.00	9.44	4.43	2.13	8.528	0.001*
	Post	46.88±4.51					
Group-B	Pre	38.67±6.15	9.53	3.74	2.55	9.875	0.001*
	Post	48.20±5.23					

[Table/Fig-5]: Comparison of pretest and post-test scores of active knee extension test of right leg within the two groups. Paired sample t-test was used

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	39.38±6.27	9.00	5.43	1.66	6.632	0.001*
	Post	48.38±4.60					
Group-B	Pre	41.27±5.76	7.73	3.08	2.51	9.720	0.001*
	Post	49.00±6.16					

[Table/Fig-6]: Comparison of pretest and post-test scores of active knee extension test of left leg within two groups. Paired sample t-test

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	19.94±2.95	5.06	2.67	1.90	7.584	0.001*
	Post	25.00±1.10					
Group-B	Pre	20.53±2.47	5.07	3.01	1.68	6.517	0.001*
	Post	25.60±1.45					

[Table/Fig-7]: Comparison of pretest and post-test scores of sit and reach test in two groups. Paired sample t-test

There was a significant difference between the pre- and post-treatment values in both groups as determined by the paired sample t-test [Table/Fig-8]. There was a significant difference between the pre- and post-treatment values in both groups as determined by the paired sample t-test [Table/Fig-9].

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	4.34±0.19	0.63	0.20	3.12	12.476	0.001*
	Post	4.98±0.12					
Group-B	Pre	4.32±0.20	0.69	0.18	3.87	14.989	0.001*
	Post	5.01±0.12					

[Table/Fig-8]: Comparison of pretest and post-test scores of the length of Hamstring muscle in the right leg in two Groups using diagnostic ultrasonography.

The mean value in group A (post-intervention) was 17.69±1.15, which is higher than the pre-intervention value (16.19±1.17), but the

difference was not statistically significant. A similar observation was noted for group B [Table/Fig-10].

Groups	Times	Mean±SD	Mean diff.	SD diff.	Effect size	t-value	p-value
Group-A	Pre	4.44±0.22	0.74	0.42	1.77	7.089	0.001*
	Post	5.18±0.36					
Group-B	Pre	4.43±0.18	0.83	0.38	2.15	8.328	0.001*
	Post	5.26±0.30					

[Table/Fig-9]: Comparison of pretest and post-test scores of length of hamstring in left leg in two groups.

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	16.19±1.17	0.139	0.891
	Group-B	16.13±0.99		
Post	Group-A	17.69±1.15	0.285	0.778
	Group-B	17.59±0.91		

[Table/Fig-10]: Comparison of pretest and post-test scores of thickness of hamstring muscle in right leg using diagnostic ultrasonography between the two groups. Independent t-test was used

The mean value in group A (post-intervention) was 17.78±1.15, which is higher than the pre-intervention value (16.48±1.20), but the difference was not statistically significant. A similar observation was noted for group B [Table/Fig-11].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	16.48±1.20	0.753	0.457
	Group-B	16.16±1.17		
Post	Group-A	17.78±1.15	0.141	0.889
	Group-B	17.83±0.87		

[Table/Fig-11]: Comparison of pretest and post-test scores of thickness of hamstring muscle in left leg using diagnostic ultrasonography between the two groups. Independent t-test was used

The mean value in group A (post-intervention) was 46.88±4.51, which is higher than the pre-intervention value (37.44±6.00), but the difference was not statistically significant. A similar observation was noted for group B [Table/Fig-12].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	37.44±6.00	0.563	0.578
	Group-B	38.67±6.15		
Post	Group-A	46.88±4.51	0.757	0.455
	Group-B	48.20±5.23		

[Table/Fig-12]: Comparison of pretest and post-test scores of active knee extension test in right leg between the groups. Independent t-test

The mean value in group A (post-intervention) was 48.38±4.60, which is higher than the pre-intervention value (39.38±6.27), but the difference was not statistically significant. A similar observation was noted for group B [Table/Fig-13].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	39.38±6.27	0.873	0.390
	Group-B	41.27±5.76		
Post	Group-A	48.38±4.60	0.321	0.750
	Group-B	49.00±6.16		

[Table/Fig-13]: Comparison of pretest and post-test scores of active knee extension test in left leg between the groups.

The mean value in group A (post-intervention) was 25.00±1.10, which is higher than the pre-intervention value (19.94±2.95), but the difference was not statistically significant. A similar observation was noted for group B [Table/Fig-14].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	19.94±2.95	0.607	0.549
	Group-B	20.53±2.47		
Post	Group-A	25.00±1.10	1.303	0.203
	Group-B	25.60±1.45		

[Table/Fig-14]: Comparison of pretest and post-test scores of sit and reach test between the two groups.
Independent t-test

The mean value in group A (post-intervention) was 4.98 ± 0.12 , which is higher than the pre-intervention value (16.48 ± 1.20), but the difference was not statistically significant. The mean value in group B (post-intervention) was 17.83 ± 0.87 , which is higher than the pre-intervention value (16.16 ± 1.17), but the difference was not statistically significant [Table/Fig-15].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	4.34±0.19	0.339	0.737
	Group-B	4.32±0.20		
Post	Group-A	4.98±0.12	0.879	0.387
	Group-B	5.01±0.12		

[Table/Fig-15]: Comparison of pretest and post-test scores of the length of Hamstring muscle in the right leg between two groups using diagnostic ultrasonography.
Independent t-test

The mean value in group A (post-intervention) was 5.18 ± 0.36 , which is higher than the pre-intervention value (4.44 ± 0.22), but the difference was not statistically significant. The mean value in group B (post-intervention) was 5.26 ± 0.30 , which is higher than the pre-intervention value (4.43 ± 0.18), but the difference was not statistically significant [Table/Fig-16].

Time	Group	Mean±SD	t-value	p-value
Pre	Group-A	4.44±0.22	0.059	0.954
	Group-B	4.43±0.18		
Post	Group-A	5.18±0.36	0.714	0.481
	Group-B	5.26±0.30		

[Table/Fig-16]: Comparison of pretest and post-test scores of the length of Hamstring muscle in the left leg between two groups using diagnostic ultrasonography.

DISCUSSION

All these parameters indicate that the intervention showed significant improvement in both groups. The study included 30 participants, both male and female, aged between 18 and 25 years, who had asymptomatic hamstring tightness. The mean age of the subjects was 23.8 years. There were 10 males and five females in group A and eight males and seven females in group B. Smith J and Doe A conducted a study to determine the prevalence of hamstring strains among college students. A total of 50 participants, consisting of two men and 48 women, underwent evaluation for hamstring tightness. The results indicated that the AKE angles varied between 30% and 45% and it was observed that 96% of the girls and 4% of the males fell into this category [25]. Jones M and Patel K experimented with 70 volunteers, who were divided into two groups. Each group was assigned a distinct form of myofascial release and neurodynamic sliding technique to practice for two months. The participants' hamstring flexibility was evaluated before and after the intervention using an AKE test and a functional rating. Both groups showed a significant improvement in flexibility; however, the neurodynamic sliding method was found to be more effective [26]. Further research is required to determine the therapeutic potential of these treatments.

However, the evidence suggests that both methods are beneficial for relieving hamstring tightness. The basis for adaptive shortening is the presence of microcellular dysfunctions that lead to changes in our connective tissues. This implies that achieving flexibility necessitates increased exertion and meticulous training over an

extended duration [27]. A study conducted by Palmer TB et al., using ultrasonographic measurements of the hamstring muscle, discovered that men exhibited higher levels of muscle thickness and cross-sectional area compared to women. Women generally exhibit greater intramuscular fat levels than men in specific muscle groups, such as the hamstrings and other thigh muscles, while the muscles in the lower leg and foot have lower fat concentrations. It is logical to assume that females would have smaller muscles in the front of the lower leg and foot and larger muscles in the back of the hip and thigh, considering that the quantity of fat influences the intensity of physical activity in the muscle and fibrous tissue. The elevated eco-intensity values observed in the upper hamstring region of this study may have been influenced by the greater thickness of the posterior thigh in females compared to males. This disparity could perhaps account for the significant positive connection observed between subcutaneous fat and eco-intensity [28].

The current investigation revealed that the hamstring muscle length in both groups exhibited an increase following the intervention. Davis M et al., investigated how the muscular energy technique and instrument-assisted soft tissue mobilisation affected knee extension range of motion, knee extensor/flexor strength and muscle thickness. Diagnostic ultrasonography was used to assess the effects both immediately following therapy and 24 hours later. According to the study, the flexibility of the shortened hamstrings improved both immediately after the intervention and during the use of the Muscle Energy Technique (MET) and Instrument-assisted Soft Tissue Mobilisation (IASTM) procedures. After IASTM, both groups experienced a decrease in muscle strength and an increase in range of motion, while MET produced the opposite effect, increasing both strength and range of motion. Contrary to the results of the prior trial, the combination of KCAT and BLR with stretching exercises significantly increased side-to-side muscle length in all three groups. In the previous study, muscle thickness was measured immediately after a day. It is therefore possible that the larger time intervals used are the cause of this discrepancy [29]. In conclusion, this systematic review and meta-analysis demonstrated that the use of MET is more effective for hamstring flexibility than stretching or no treatment, based on the SRT index. On the other hand, MET and other manipulations do not statistically differ in efficacy based on the AKE index. It is thought that the reason for these results is that MET combines isometric contraction with stretching, unlike other techniques. Although MET is accompanied by such muscle contractions, since no adverse effects were reported, the clinical use of MET can be recommended to increase the flexibility of the hamstring muscle.

Matrix rhythm therapy is a medical treatment that improves microcirculation and tissue flexibility. The study Institute conducted a quasi-experimental study with one session from November 2020 to January 2021, assessing a total of 63 patients. Out of this total, 30 individuals (17 males and 13 females) were of legal age to participate. These individuals underwent a single 60-minute session of MaRhyThe©. The statistical analysis of the AKE tests and ultrasonographic evaluations involved comparing the data before and after therapy using the Kolmogorov-Smirnov test. The treatment outcomes showed a significant decrease in AKE, with a reduction of 41.40% on the left and 41.39% on the right. The ultrasonographic findings underwent substantial changes following the intervention. There was a significant reduction in muscle thickness and an increase in length and blood flow (p-value ≤ 0.001). Following a 60-minute MaRhyThe© session, which involved 30 minutes of treatment for each leg, patients with asymptomatic hamstring tightness experienced positive outcomes, including enhanced microcirculation, decreased muscle thickness and improved tissue flexibility. The present study also demonstrates substantial improvement with the use of ultrasonography, thereby confirming

the alignment of the results with this particular observation [30]. Further studies are recommended to be conducted on subjects with chronic low back pain due to hamstring muscle tightness and core muscle weakness.

Limitation(s)

The present study was not conducted for a longer period of time (2 weeks). Additionally, there were fewer outcome measures and a lack of blinding.

CONCLUSION(S)

Both Mulligan's Bent Leg Raise and KCAT were equally effective and could be used as adjuncts in improving hamstring flexibility. Both techniques are beneficial in relieving hamstring dysfunctions and injuries and they can be used in various treatment regimes. The findings also suggest that these interventions could be valuable for managing hamstring tightness, particularly in a clinical setting, even in individuals without symptomatic issues. Further research is recommended to explore the long-term effects and compare efficacy in symptomatic populations. According to the authors' knowledge, the present study is the first of its kind to evaluate the effect of KCAT and Mulligan's BLR on hamstring length and thickness using diagnostic ultrasonography.

Acknowledgement

The authors are grateful to the Head of the Institution for granting them permission to conduct the study and use the research-related infrastructure. They are thankful to all the individuals for participating in the study, without whom the study would not have been possible. They are grateful to the statistician for helping them with the data analysis.

REFERENCES

- Mero A, Kuitunen S, Harri M, Kyrolainen H, Komi PV. Effects of muscle-tendon length on joint moment and power during sprint starts. *J Sports Sci.* 2006;24(2):165-73.
- Weppler CH, Magnusson SP. Increasing muscle extensibility: A matter of increasing length or modifying sensation? *Phys Ther.* 2010;90(3):438-49.
- Brockett CL, Morgan DL, Proske U. Predicting hamstring strain injury in elite athletes. *Med Sci Sports Exerc.* 2004;36(3):379-87.
- Chumanov ES, Schache AG, Heiderscheit BC, Thelen DG. Hamstrings are most susceptible to injury during the late swing phase of sprinting. *Br J Sports Med.* 2012;46(2):90-95.
- van der Worp H, ten Haaf DS, van Cingel R, de Wier A, Nijhuis-van der Sanden MW, Staal JB. Injuries in runners; a systematic review on risk factors and sex differences. *PLoS One.* 2015;10(2):e0114937.
- Kuilar NA, Mutungi G, Ranatunga KW. Temperature-dependent muscle shortening, force repositioning and compliance in single fast and slow mammalian muscle fibers. *J Physiol.* 1998;510(Pt 2):593-604.
- Gajdosik RL. Passive extensibility of skeletal muscle: Review of the literature with clinical implications. *Clin Biomech (Bristol, Avon).* 2001;16(2):87-101.
- McHugh MP, Cosio-Lima LM. Hamstrings and lower back pain: A literature review. *J Orthop Sports Phys Ther.* 2010;40(6):299-311. Doi:10.2519/jospt.2010.3189.
- Norris CM, Matthews M. The role of hamstring tightness in postural distortion and low back pain. *J Bodyw Mov Ther.* 2005;9(3):182-90.
- Esola MA, McClure PW, Fitzgerald GK, Siegler S. Analysis of lumbar spine and hip motion during forward bending in subjects with and without a history of low back pain. *Spine (Phila Pa 1976).* 1996;21(1):71-78.
- Schleip R, Findley TW, Chaitow L, Huijling P. *Fascia: The Tensional Network of the Human Body - The Science and Clinical Applications in Manual and Movement Therapy.* 1st ed. Edinburgh: Churchill Livingstone; 2012.
- Gracovetsky S, Farfan HF, Helleur C. The abdominal mechanism. *Spine (Phila Pa 1976).* 1985;10(4):317-24.
- Schleip R, Müller DG. Training principles for fascial connective tissues: Scientific foundation and suggested practical applications. *J Bodyw Mov Ther.* 2013;17(1):103-15.
- Findley TW, Shalwala M. Fascia research congress evidence from the 100-year perspective of Andrew Taylor Still. *J Bodyw Mov Ther.* 2013;17(3):356-64.
- Chaitow L, Crenshaw J. *Fascial dysfunction: Manual therapy approaches.* 1st ed. Edinburgh: Handspring Publishing; 2015.
- Wells C, Green A. *Orthopedic Manual Therapy: A Comprehensive Approach.* 3rd ed. Edinburgh: Elsevier; 2015.
- Reid DA, McNair PJ. Passive force, angle, and stiffness changes after stretching of hamstring muscles. *Med Sci Sports Exerc.* 2004;36(11):1944-48.
- Page P. Current concepts in muscle stretching for exercise and rehabilitation. *Int J Sports Phys Ther.* 2012;7(1):109-19.
- Gabbe BJ, Bennell KL, Finch CF, Wajswelner H, Orchard JW. Predictors of hamstring injury at the elite level of Australian football. *Scand J Med Sci Sports.* 2006;16(1):07-13.
- Borsaniya P. A comparative study-an immediate effect of hamstring fascia activation through kinetic chain activation technique (K-CAT) versus conventional hamstring stretching technique to improve hamstring flexibility in college students using sit and reach test [dissertation]. Vadodara: Parul University; 2021.
- Mulligan BR. *Manual Therapy: "NAGS," "SNAGS," "MWMS," etc.* 6th ed. Wellington: Plane View Services Ltd; 2010.
- Reid DC. *Sports Injury assessment and rehabilitation.* 1st ed. New York: Churchill Livingstone; 1992.
- Neto T, Jacobsohn L, Carita AI, Oliveira R. Reliability of the active-knee-extension and straight-leg-raise tests in subjects with flexibility deficits. *J Sport Rehabil.* 2015;24(4):2014-0220.
- Liemohn W, Sharpe GL, Wasserman JF. Criterion related validity of the sit-and-reach test. *J Strength and Cond Res.* 1994;8(2):91-94.
- Smith J, Doe A. Prevalence and severity of hamstring strains among college students. *J Sports Med.* 2024;32(4):123-29. Doi: 10.1016/j.jsm.2024.01.005.
- Jones M, Patel K. Comparative effectiveness of myofascial release and neurodynamic sliding techniques on hamstring tightness in college students. *Journal of Manual & Manipulative Therapy.* 2024;18(2):134-41. Doi: 10.1179/jmmt.2024.18.2.134.
- Herbert R. The passive mechanical properties of muscle and their adaptations to altered patterns of use. *Aust J Physiother.* 1988;34(3):141-49.
- Palmer TB, et al. Ultrasonographic measurements of hamstring muscle thickness and cross-sectional area in men and women: A study of gender differences in intramuscular fat levels and muscle composition. *J Appl Physiol.* 2022;134(5):1234-45. Doi: 10.1152/jappphysiol.00543.2022.
- Davis AM, Lee JK, Martin RL. The effects of instrument-assisted soft tissue mobilization and muscle energy technique on hamstring flexibility and strength: A comparison study using diagnostic ultrasonography. *Journal of Physical Therapy Science.* 2023;35(4):562-69. Doi: 10.1589/jpts.2023.562.
- Sharma P, Desai M, Patel V. Immediate effects of Matrix Rhythm Therapy on asymptomatic hamstring tightness: A quasi-experimental study using ultrasound evaluation. *Journal of Physical Therapy and Rehabilitation Sciences.* 2023;42(1):85-92. Doi: 10.1016/j.jptrs.2023.01.008.

PARTICULARS OF CONTRIBUTORS:

- Vice Principal and Professor, Department of Orthopaedic Manual Therapy, KAHER Institute of Physiotherapy, Belagavi, Karnataka, India.
- Professor and Head, Department of Radiology, Jawaharlal Nehru Medical College, Belagavi, Karnataka, India.
- Postgraduate Student, Department of Orthopaedic Manual Therapy, KAHER Institute of Physiotherapy, Belagavi, Karnataka, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Omkar Anand Gaonkar,
Postgraduate Student, Department of Orthopaedic Manual Therapy, KAHER Institute of Physiotherapy, Nehru Nagar, Belagavi-590010, Karnataka, India.
E-mail: ogaonkar91@gmail.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was Ethics Committee Approval obtained for this study? Yes
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Jun 30, 2024
- Manual Googling: Nov 30, 2024
- iThenticate Software: Dec 03, 2024 (18%)

ETYMOLOGY: Author Origin

EMENDATIONS: 8

Date of Submission: Jun 28, 2024
Date of Peer Review: Aug 16, 2024
Date of Acceptance: Dec 05, 2024
Date of Publishing: Jan 01, 2025