

Microabrasion as an Effective Treatment for Severe Dental Fluorosis: Images in Medicine

PRIYANKA RAJESH BHOJWANI¹, ANUJA IKHAR², ADITYA PATEL³, MANOJ CHANDAK⁴, SHWETA SEDANI⁵

(CC) BY-NC-ND

Keywords: Hypomineralisation, Hydrochloric acid, Pumice slurry, Phosphoric acid

A 28-year-old male presented to the dental clinic with white and brown discolourations, as well as hard, pitted areas on the enamel [Table/Fig-1,2], seeking aesthetic treatment. After obtaining a detailed medical history and performing a clinical examination, including electrical pulp testing and cold testing, a radiographic evaluation was conducted. The radiographs revealed no periapical lesions or caries, and no signs of resorption were observed in the teeth of interest. All the associated teeth were vital, with a healthy periodontium. The patient was diagnosed with severe enamel fluorosis, classified as grade 4 according to the Dean's Index [1].

Treatment options, including veneers, crowns, and macroabrasion, were discussed with the patient and his parents. Ultimately, the enamel microabrasion technique was selected to address the stains and pitted areas on the facial surfaces of the upper incisors, canines, and premolars.

After obtaining informed consent for both the treatment and the publication of identifiable images, petroleum jelly was applied to protect the gingival tissues. The operative field was isolated using a rubber dam to shield the gums from the microabrasive compound. All individuals involved in the procedure, including the patient, assistants, and operator, wore appropriate personal protective equipment and eye protection throughout the treatment.

For the enamel microabrasion procedure, 6.6% hydrochloric acid was freshly prepared in the biochemistry laboratory. The acid was applied using a rubber cup attached to a low-speed handpiece, with firm pressure applied to remove the stains. Pumice was selected as the abrasive for this case. The compound was applied three times, with each application lasting one minute for six teeth-approximately 10s per tooth- followed by water rinsing between each session. Afterward, the abraded enamel surface was polished with a fluoridated paste, and a neutral 2% sodium fluoride gel was applied for four minutes. [Table/Fig-3] shows the post-treatment appearance of the enamel following the microabrasion procedure.

One month later, dental bleaching was performed using 35% hydrogen peroxide (SDI Pola Office Bleaching Material) with the aid of an Optradam. A single 20-minute cycle was performed. To prevent reversal of the process, a polishing kit was used in sequence. Immediate postoperative images are shown in [Table/Fig-4,5].

Telephonic follow-up was conducted for three days. The patient reported mild sensitivity only on the day of the procedure, immediately following the treatment, which resolved within four hours. No other side-effects or complications were reported during or after the treatment. Three months later, the patient was recalled for a follow-up visit. During this visit, the pits were restored with composite (Dentsply Spectrum Composite, Shade A1). The follow-up image is shown in [Table/Fig-6]. [Table/Fig-7] presents a comparison of the pre-operative and postoperative images.

The results showed a significant improvement in colour, with the shade lightening from A2 to A1. According to the Dean's Modified



[Table/Fig-1]: Pre-operative clinical images.



[Table/Fig-2]: Pre-operative radiographic images.



[Table/Fig-3]: Postoperative clinical images post-microabrasion.

Index, the postoperative image indicated a very mild result, corresponding to grade 1 [2].

The microabrasion process, which involves acidic agents and abrasive compounds, helps reduce the whitish appearance of lesions by removing the outer dysplastic enamel layer. A study by Lu Y et al., divided participants into two groups for office bleaching, one of which combined microabrasion using 37% phosphoric acid. The results revealed a significant difference, with the microabrasion group showing better outcomes [3]. Similarly, Divyameena B et al., achieved



[Table/Fig-4]: Postoperative clinical image under rubber dam after bleaching.



[Table/Fig-5]: Postoperative clinical images post bleaching.



[Table/Fig-6]: Follow-up clinical images after composite restoration of the pits.

comparable results using Opalustre for microabrasion and observed no postoperative sensitivity [4]. However, according to a systematic review, microabrasion is considered effective for mild to moderate cases but shows less promising results for more severe cases [5].

A newer treatment modality that has gained acceptance is resin infiltration. In a study by Reddy VN et al., where 37% phosphoric acid with pumice, Opalustre, and resin infiltration were compared, it



[Table/Fig-7]: Pre-operative and postoperative clinical images.

was concluded that the mean surface roughness was 10.42 for the combination, while resin infiltration had a surface roughness of only 1.99. This suggests that resin infiltration is a less invasive method with reduced post-operative reversal of results [6].

This case demonstrates that microabrasion is an effective and conservative treatment option for severe dental fluorosis, offering substantial aesthetic improvements without the need for invasive procedures. By selectively removing discoloured and hypomineralised enamel layers, microabrasion enhanced the patient's smile, resulting in a more uniform and natural appearance of the teeth.

REFERENCES

- [1] Dean HT. Classification of mottled enamel diagnosis. *J Am Dent Assoc.* 1934;21(8):1421-26.
- [2] Dean HT. The oral hygiene index. *J Am Dent Assoc.* 1942;28(6):929-33.
- [3] Lu Y, Wang J, Liang Y, Chang S, Gao R. Enamel microabrasion combined with in-office bleaching in treating with moderate to severe dental fluorosis-a randomized clinical trial. *Biomed J Sci Tech Res.* 2021;34(2):26490-96.
- [4] Divyameena B, Sherwood A, Rathna Priyanga S, Deepika G. Clinical performance of enamel microabrasion for esthetic management of stained dental fluorosis teeth. *Oper Dent.* 2021;46(1):15-24. Doi: 10.2341/19-248-C. PMID: 32926733.
- [5] Blanchet I, Camoin A, Tardieu C, Jacquot B. Microabrasion in the management of enamel discolourations in paediatric dentistry: A systematic review. *J Clin Pediatr Dent.* 2023;47(1):17-26. Doi: 10.22514/jocpd.2022.015. Epub 2023 Jan 3. PMID: 36627216.
- [6] Reddy VN, Nagar P, Lakhotia R, Jenny A, Ravi PH, Kanugondappa R. Comparative evaluation of three different microabrasion techniques in esthetic management of fluorosis: An in vivo study. *Int J Clin Ped Dent.* 2023;16(4):572.

PARTICULARS OF CONTRIBUTORS:

1. Postgraduate Student, Department of Conservative Dentistry and Endodontics, Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India.
2. Associate Professor, Department of Conservative Dentistry and Endodontics, Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India.
3. Professor and Head, Department of Conservative Dentistry and Endodontics, Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India.
4. Professor and Dean, Department of Conservative Dentistry and Endodontics, Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India.
5. Associate Professor, Department of Conservative Dentistry and Endodontics, Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and Research, Wardha, Maharashtra, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Priyanka Rajesh Bhojwani,
Postgraduate Student, Department of Conservative Dentistry and Endodontics,
Sharad Pawar Dental College, Datta Meghe Institute of Higher Education and
Research, Wardha-442001, Maharashtra, India.
E-mail: pbhojwani0000@gmail.com

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Jan 10, 2025
- Manual Googling: Mar 12, 2025
- iThenticate Software: Mar 29, 2025 (13%)

ETYMOLOGY: Author Origin

EMENDATIONS: 7

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

Date of Submission: Jan 07, 2025

Date of Peer Review: Feb 24, 2025

Date of Acceptance: Apr 02, 2025

Date of Publishing: Jul 01, 2025