

A Narrative Review of Different Nasotracheal Intubation Techniques with Recent Advancements in Clinical Practice

YAN-QIU ZHANG¹, JI ZHAO², CHENG-CHENG LIANG³, YONG WANG⁴

ABSTRACT

Nasotracheal intubation is commonly used in clinical practice for procedures requiring general anaesthesia, mechanical ventilation, and emergency treatment of critically ill patients. It presents a new good surgical field and operating space for maxillofacial surgeries, offers an artificial airway for long-term mechanically ventilated patients in the Intensive Care Unit (ICU), and provides a method for emergency and difficult airway intubations. Compared to orotracheal intubations, nasotracheal intubation has advantages such as minimal stimulation to the throat, good patient tolerance, easy fixation, facilitation of oral care, and longer duration of placement. However, it requires higher level of technical skill from the operator and carries a higher incidence of complications like epistaxis and sinusitis, which have restricted its widespread application. From the early blind intubation method to the current direct laryngoscopy, video laryngoscopy, and fiberoptic bronchoscopy, the success rate of nasotracheal intubation has increased, while the incidence of complications has decreased. This review summarises the common methods, advantages and disadvantages of nasotracheal intubation, as well as the research progress on related complications and coping strategies, aiming to enhance the application of this technology in clinical practice.

Keywords: Complications, Epistaxis, Fiberoptic bronchoscopy, Nasal pressure injury, Sinusitis, Tracheal intubation, Video laryngoscopy

INTRODUCTION

Tracheal intubation is a routine procedure for rescuing critically ill patients and for use under general anaesthesia, with the main purpose of maintaining a clear airway, preventing reflux and aspiration, as well as connecting to a ventilator for mechanical ventilation. Tracheal intubation can be performed by the oral or the nasal route. Orotracheal intubation is characterised by a simple and convenient operation, but patients may find it uncomfortable, and it is not conducive to oral care. Nasotracheal intubation is more in line with the physiological characteristics of the patient's airway, with advantages such as minimal stimulation to the pharynx, good patient tolerance, easy fixation, convenient oral care, and longer indwelling time. However, it requires a high level of technical expertise from the operator and carries risks of nasal and nasopharyngeal injury, epistaxis, and sinusitis. In recent years, with the development of clinical medicine and advancements in research, especially the application of technologies such as video laryngoscopy and fiberoptic bronchoscopy in nasotracheal intubation, the success rate of nasotracheal intubation has significantly improved while lowering associated complications [1]. This article provides a comprehensive review of the research progress on the application of nasotracheal intubation in clinical practice.

Nasotracheal intubation is a frequently utilised technique in patients undergoing oral and maxillofacial surgery, offering improved surgical visibility and operational space compared with orotracheal intubation [2,3]. For anterior cervical surgery, nasotracheal intubation obviates the need for excessive neck extension, with the mandibular-cervical angle increased by 7.3°, thereby improving the surgical field [4]. In the ICU, tracheal intubation is predominantly conducted through the oral route, as opposed to the nasotracheal approach. This preference is largely attributable to complications associated with nasotracheal intubation, including epistaxis and infections such as sinusitis and ventilator-associated pneumonia. Consequently, orotracheal intubation is increasingly being utilised as a replacement for nasotracheal intubation [5].

However, in the ICU, patients requiring prolonged airway management are common, prognosis is often poor, and family members are often unwilling to let them undergo tracheostomy. Nasotracheal intubation can be considered for such patients as the duration of orotracheal intubation should not be prolonged. For some awake patients in the ICU, nasotracheal intubation can reduce the dosage of sedatives, and some patients can eat on their own without the need for a feeding tube. Furthermore, in patients with respiratory infectious diseases, nasotracheal intubation can reduce the risk of infection for healthcare workers during intubation. In the Paediatric ICU (PICU), the proportion of nasotracheal intubation seems to be higher than in adult patients, to reported rates in PICU children being 5.6% and 3.8%, respectively [6,7]. A study in the context of cardiac surgery indicated that the proportion of neonates undergoing nasotracheal intubation was 41%, while the corresponding figure for infants was 38%. These findings reinforce the notion that nasotracheal intubation in paediatric patients is safe and does not significantly elevate the risk of severe complications. Compared with orotracheal intubation, there is a marked reduction in the rate of unplanned extubation, and the incidence of sinusitis and ventilator-associated pneumonia remains statistically unchanged [8].

It is crucial to note that bilateral nasal obstruction constitutes an absolute contraindication for nasotracheal intubation. Therefore, a thorough assessment of the nasal cavity's patency should be conducted prior to the procedure [6]. Additionally, certain conditions, such as nasopharyngeal haemangioma, severe laryngeal oedema, coagulation disorders, skull base fractures, and the possibility of elevated intracranial pressure, are considered relative contraindications. Patients presenting with these conditions should avoid nasotracheal intubation whenever possible.


The Choice of Nostril

When nasotracheal intubation is performed, if both nasal cavities are clear, intubation can be done through either nostril. A study reported that the success rates of intubation through the left nostril and the right nostril were similar {Group L 96% (24/25), Group R 96% (24/25)},

with no difference in the incidence and severity of epistaxis during intubation [9]. However, the most current available evidence suggests that intubation through the right nostril is safer and faster, with a lower incidence and severity of epistaxis [10,11]. Research has shown that for intubation through the left nostril, angling the tip of the endotracheal tube towards the nasal septum can reduce the incidence of epistaxis [12]. A study of 390 patients undergoing nasotracheal intubation found that among 94 patients with nasal septum deviation identified by X-ray, complications occurred more frequently when nasal septum deviation was present (35.0% vs 18.5%) [13].

Nasotracheal Intubation Methods

Nasotracheal intubation does not require specialised tracheal tubes, but opting for a smaller endotracheal tube compared with oral intubation enhances ease of use and reduces complications. The Parker Flex-Tip™ tracheal tube, featuring a soft, flexible tip, minimises nasal mucosal damage during insertion [Table/Fig-1] [14]. Additionally, it helps regulate haemodynamic responses, with minimal effects on heart rate and diastolic pressure [15]. Preheating and softening the tube prior to intubation can further reduce nasal mucosal injury and lower the incidence of epistaxis [16]. A study on 140 patients found significantly reduced epistaxis severity in the heat-softened group (7% vs. 51%) compared to the control group



[Table/Fig-1]: Tip difference between conventional-tip and Parkertip tracheal tubes: a) Top view; b) Lateral view [14].
Image source: [14]

[17]. However, excessive softening may compromise tracheal tube patency, potentially affecting intubation success.

Blind Nasotracheal Intubation (BNTI): Blind probing was initially utilised for nasotracheal intubation due to equipment constraints, without any auxiliary measures, requiring a high level of technical proficiency from the operator. Trained operators can also achieve a higher success rate. In 1993, van Elstraete AC et al., reported a success rate of 75% for blind nasotracheal intubation using cuff-inflation method, with an overall success rate of 95% [18]. The specific procedure involved inserting the endotracheal tube through the right nostril, positioning the tube tip at the oropharynx, inflating the cuff with 15 mL of gas, gently advancing the tube until encountering slight resistance when the cuff touched the vocal cords, deflating the cuff, and inserting the endotracheal tube into the trachea. Another method is the suction catheter-guided blind nasal intubation, where a suction catheter is first inserted through the nasal cavity into the trachea, followed by sliding the endotracheal tube over the suction catheter and slowly advancing it into the trachea before removing the suction catheter [19,20]. Training operator providers in blind nasotracheal intubation is crucial, with proficient individuals achieving success rates comparable to fiberoptic bronchoscope nasotracheal intubation [21]. Despite being considered outdated, this technique remains a critical option when dealing with special patients such as severe cervical deformities, burns, and those with limited or impossible mouth opening [22].

Direct Laryngoscopy Nasotracheal Intubation (DLNTI): Direct laryngoscopy has been the standard method for tracheal intubation since its development in the early 20th century. It enables visualisation of the vocal cords, improving intubation success over blind techniques. Magill forceps assist in guiding the tube into the trachea, reducing difficulty and increasing success rates. Using a disposable tube core can replace Magill forceps, significantly reducing intubation time in patients without difficult airways [23]. However, DLNTI requires sedation and anaesthesia, poses cardiovascular risks, and is gradually being replaced by video laryngoscopy [24].

Video Laryngoscopy Nasotracheal Intubation (VLNTI): Video laryngoscopy has improved intubation success rates and efficiency. Unlike direct laryngoscopy, it does not require excessive head extension, making positioning easier [25]. In ICU and emergency settings, it enhances intubation success and reduces procedural difficulty [26,27]. It also improves first-pass success and lowers complication rates compared to direct laryngoscopy [28,29]. Paediatric patients benefit from HugeMed® video laryngoscopy, which enhances vocal cord visualisation and minimises tissue trauma [30]. Studies confirm faster intubation times and reduced use of Magill forceps with video laryngoscopy [31]. McGrath and C-MAC® video laryngoscopes provide superior vocal cord visualisation, reduced airway trauma, and stable haemodynamics [32,33]. C-MAC's adjustable blade angle aids nasotracheal intubation in cervical injury patients [34]. For awake intubation, video laryngoscopy is faster and safer than fiberoptic bronchoscopy, reducing oxygen desaturation risks [35].

Fiberoptic Bronchoscope Nasotracheal Intubation (FBNTI): FBNTI involves inserting a fiberoptic bronchoscope through the nasal cavity to guide the tracheal tube into the trachea. This technique offers direct visualisation, ensuring proper tube placement and minimising misplacement risks. It significantly lowers epistaxis incidence of epistaxis compared with blind techniques and direct laryngoscopy intubation [36]. In paediatric dental procedures, FBNTI is safe and yields higher success rates with fewer complications than direct laryngoscopy [37]. Traditional bronchoscopes are rigid, but flexible models reduce complications, especially in awake patients [38]. Oral secretions and bleeding can hinder success, requiring mitigation efforts [39]. Despite its advantages, FBNTI involves extensive preparation, limiting its use in emergency intubation. Each technique has specific benefits and limitations, with video

laryngoscopy emerging as the preferred method in various clinical settings.

Other Nasotracheal Intubation Methods

Lightwand-guided nasotracheal intubation: This method utilises a light source at the tip of a flexible lightwand to guide tracheal tube placement. Compared blind intubation, it offers higher success rates, fewer complications, and shorter intubation times [40]. The process involves inserting the tracheal tube through the nose into the oropharynx, followed by inserting the lightwand into the tube. The light's positioning just above the cricoid cartilage helps ensure proper placement before advancing the tube into the trachea and removing the lightwand. However, repeated adjustments may risk injury to the nasal cavity, epiglottis, or airway.

Ultrasound-Guided Tracheal Intubation (UGTI): UGTI is a newer approach beneficial for patients with restricted mouth opening who cannot undergo laryngoscopy. Unlike traditional methods, it is unaffected by oral secretions or bleeding [41]. While widely applied for orotracheal intubation, reports on its nasotracheal use remain limited.

Comparison of Intubation Methods and Process Flowchart

Various nasotracheal intubation methods are used clinically, each with its advantages, disadvantages, and indications. Operators should select the most appropriate technique based on patient-specific factors. A comparative analysis of these methods is provided in [Table/Fig-2] [18,22,24,26,28-34,36-38], while the step-by-step nasotracheal intubation procedure is outlined in [Table/Fig-3] [1].


Intubation methods	Advantages	Disadvantages	Success rates	Optimal indications
BNTI	No special equipment required [18]	High failure rate [18]	Low [18]	Severe cervical deformities; burns, and patients with limited or impossible mouth opening [22]
DLNTI	Fast speed [24]	Need sedation and analgesia	High [24]	Patients without difficult airways [24]
VLNTI	Easy to operate, fast speed [26,28-32,34]	Need video laryngoscope	High [26,28-32,34]	Widely used (Emergency, Anaesthesiology, ICU) [26,28,29,31-34]
FBNTI	High success rate [37,38]	Cumbersome preparation before intubation [36,37,38]	High [37,38]	Surgery patients undergoing preoperative anaesthesia intubation [37,38]

[Table/Fig-2]: Comparative analysis of various intubation methods [18,22,24,26,28-34, 36-38].
BNTI: Blind nasotracheal intubation; DLNTI: Direct laryngoscopy nasotracheal intubation; VLNTI: Video laryngoscopy nasotracheal intubation; FBNTI: Fiberoptic bronchoscope nasotracheal intubation

Complications of Nasotracheal Intubation

Epistaxis: Epistaxis, a frequent and severe complication of nasotracheal intubation, results from damage to the nasal mucosa and vascular damage. Excessive bleeding can be life-threatening, necessitating preventive measures. Guidelines recommend local nasal vasoconstrictors pre-intubation [42]. Pre-emptive epinephrine nebulisation and xylometazoline spray effectively reduce epistaxis rates [43,44]. Xylometazoline drops have been shown to further decrease severe epistaxis during intubation and extubation [45]. Studies indicate that cocaine and xylometazoline offer comparable efficacy in preventing epistaxis [46]. Nasal packing with bupivacaine effectively minimises bleeding [47], while softer North Polar Tubes further lower the epistaxis incidence [48]. Gentle intubation technique and careful tube adjustments are crucial for prevention.

Nasal pressure injury: Nasal pressure injuries are common after nasotracheal intubation but can be mitigated with protective strategies. Hydroactive dressings significantly reduce nasal wing skin



[Table/Fig-3]: Flowchart of nasotracheal intubation steps.

damage [49,50], while hydrocolloid dressings help paediatric patients by reducing pressure and absorbing exudates [51]. A 3M Microfoam™ surgical tape offers effective nasal protection [52].

Sinusitis: Prolonged nasotracheal intubation increases the risk of sinusitis due to mucosal damage and ostial obstruction. Patients intubated for over five days face a higher likelihood of developing sinusitis [53]. Late-onset sinusitis is more common with nasotracheal versus oral intubation [54]. 2% nasal mupirocin ointment may lower infection rates by reducing Staphylococcus aureus colonisation [55].

Other complications: Postoperative sore throat is prevalent, affecting 74.6% of patients following oral and maxillofacial surgery [56]. Fiberoptic bronchoscope intubation reduces sore throat compared with Macintosh laryngoscopy [57]. Cardiac arrest, triggered by nasal mucosal stimulation and the rhino-cardiac reflex, requires immediate intubation cessation and emergency drug intervention with atropine or epinephrine [58]. Rare complications include internal carotid artery injury, submucosal retropharyngeal dissection, pyriform sinus perforation, and cervicothoracic emphysema.

Given the potential severity of these complications, careful patient assessment is essential before intubation. Gentle handling minimises epistaxis risk, proper nasal protection prevents pressure injuries, and timely extubation mitigates sinusitis. Close monitoring during intubation ensures early detection of serious complications such as cardiac arrest.

Techniques to Improve the Success Rate of Nasotracheal Intubation

The primary concerns with nasotracheal intubation are success rates and complications. A study of 86 patients showed that the nasal tip-lifting method significantly improved the success rate (79.1% vs 51.2%). This technique involves selecting the appropriate nostril, lifting the nasal tip upwards with one hand, and inserting the tracheal tube with the other [59].

While the incidence of epistaxis was similar in both groups (18.6% vs 32.6%), nasal tip lifting helps the tube pass more smoothly through the nasal cavity. Research indicates that directing the tube toward the patient's head rather than the left side reduces nasal bleeding during jaw surgery [60]. Additionally, expanding the nasal cavity before intubation aids in guiding the tube into the pharynx and minimises trauma and epistaxis. Placing a nasopharyngeal airway beforehand increases stability and space, making tube insertion easier.

Finally, standardised training, selecting appropriate intubation methods, and avoiding excessive force are essential for improving intubation success and reducing complications.

CONCLUSION(S)

In clinical practice, nasotracheal intubation is a valuable airway management technique that offers advantages in specific situations. Operators must have a comprehensive understanding of indications, complications, and the latest advances in airway management to ensure the safety and effectiveness. Various nasotracheal intubation methods have their pros and cons, and healthcare providers should be proficient in one or more techniques to perform nasotracheal intubation successfully when necessary, ensuring patient safety and meeting surgical requirements.

REFERENCES

- [1] Park DH, Lee GA, Jeong CY, Yang HS. Nasotracheal intubation for airway management during anaesthesia. *Anaesth Pain Med*. 2021;16(3):232-47.
- [2] Kumar A, Gupta N, Bhargava T, Gupta A, Kumar V, Bharti SJ, et al. A comparative evaluation of fiberoptic bronchoscopy versus C-MAC D-BLADE-guided videolaryngoscopy for nasotracheal intubation under general anaesthesia in oropharyngeal carcinoma surgery patients. *Can J Anaesth*. 2024;71(4):503-10.
- [3] Ketata H, Zouche I, Sellami M, Ghars O, Ketata S, Charfeddine I, et al. Medcaptain® video laryngoscope improves nasotracheal intubation in maxillofacial surgery: A randomized clinical trial. *Trends Anaesth Crit Care*. 2023;52:101287.
- [4] Kang JH, Im SB, Jeong JH, Lee SJ, Cho SH, Hossain M. Effectiveness of nasotracheal intubation in anterior cervical surgery including C3 lesions. *Br J Neurosurg*. 2024;38(1):111-15.
- [5] Hariri G, Baudel JL, Dubee V, Dumas G, Joffre J, Bourcier S, et al. Nasotracheal intubation in ICU: An unfairly forgotten procedure. *Minerva Anesthesiol*. 2018;84(8):997-98.
- [6] Christian CE, Thompson NE, Wakeham MK. Use and outcomes of nasotracheal intubation among patients requiring mechanical ventilation across U.S. PICUs. *Pediatric Crit Care Med*. 2020;21(7):620-24.
- [7] Ducharme-Crevier L, Furlong-Dillard J, Jung P, Chiusolo F, Malone MP, Ambati S, et al. Safety of primary nasotracheal intubation in the pediatric intensive care unit (PICU). *Intensive Care Med Paediat. Neonatal*. 2024;2(1):7.
- [8] Greene NH, Jooste EH, Thibault DP, Wallace AS, Wang A, Vener DF, et al. A study of practice behavior for endotracheal intubation site for children with congenital heart disease undergoing surgery: Impact of endotracheal intubation site on perioperative outcomes-an analysis of the society of thoracic surgeons congenital cardiac anaesthesia society database. *Anaesth Analg*. 2019;129(4):1061-68.
- [9] Wang LK, Zhang X, Wu HY, Cheng T, Xiong GL, Yang XD. Impact of choice of nostril on nasotracheal intubation when using video rigid stylet: A randomized clinical trial. *BMC Anaesthesiol*. 2022;22(1):360.
- [10] Arslan ZI, Türkyilmaz N. Which nostril should be used for nasotracheal intubation with Airtraq NT®: The right or left? A randomized clinical trial. *Turk J Med Sci*. 2019;49(1):116-22.
- [11] Tan YL, Wu ZH, Zhao BJ, Ni YH, Dong YC. For nasotracheal intubation, which nostril results in less epistaxis: Right or left? A systematic review and meta-analysis. *Eur J Anaesthesiol*. 2021;38(11):1180-86.
- [12] Park JY, Yu J, Kim CS, Mun T, Jeong WS, Choi JW, et al. Reverse tube direction and epistaxis in left nasotracheal intubation: A randomized controlled trial. *Korean J Anaesthesiol*. 2024;77(6):596-604.
- [13] Chi SI, Park S, Joo LA, Shin TJ, Kim HJ, Seo KS. Identifying the more suitable nostril for nasotracheal intubation using radiographs. *J Dent Anaesth Pain Med*. 2016;16(2):103-09.
- [14] Sanuki T, Hirokane M, Matsuda Y, Sugioka S, Kotani J. The Parker Flex-Tip™ tube for nasotracheal intubation: The influence on nasal mucosal trauma. *Anaesthesia*. 2010;65(1):08-11.
- [15] Bansal T, Singhal S, Aggarwal A. A study to evaluate and compare Parker Flex-Tip tube and polyvinyl chloride endotracheal tube for hemodynamic response during intubation. *The Indian Anaesthetists' Forum*. 2024;25(1):35-41.
- [16] Prashant HT, Kerai S, Saxena KN, Wadhwa B, Gaba P. Comparison of cuff inflation method with curvature control modification in thermosoftened endotracheal tubes during nasotracheal intubation-A prospective randomised controlled study. *Indian J Anaesth*. 2021;65(5):369-76.
- [17] Kim EM, Chung MH, Lee MH, Choi EM, Jun IJ, Yun TH, et al. Is tube thermosoftening helpful for videolaryngoscope-guided nasotracheal intubation? A randomized controlled trial. *Anaesth Analg*. 2019;129(3):812-18.
- [18] van Elstraete AC, Pennant JH, Gajrai NM, Victory RA. Tracheal tube cuff inflation as an aid to blind nasotracheal intubation. *Br J Anaesth*. 1993;70(6):691-93.
- [19] Song J, Pyeon T, Kim D, Lee C, Kim J. Endobronchial suction catheter guided nasotracheal intubation with preformed nasotracheal tube. *Med Biol Sci Eng*. 2020;3(1):20-23.
- [20] Kim J, Jang EA, Kang D, Moon S, Bae HB. Comparison of the nasal cavity guidance methods' effects during nasotracheal intubation using a preformed nasotracheal tube: A prospective randomized controlled trial. *Int J Environ Res Public Health*. 2023;20(5):4503.
- [21] Parkey S, Erickson T, Hayden EM, Brown III CA, Carlson JN. Flexible nasotracheal intubation compared to blind nasotracheal intubation in the setting of simulated angioedema. *Am J Emerg Med*. 2019;37(1):1995-98.
- [22] Brezic N, Gligorevic S, Stanisavljevic J, Hadzibegovic A, Jovanovic B. Blind nasotracheal intubation in a patient with ankylosing spondylitis and severe cervical spine deformities: A case report on challenges and approaches to difficult airway management. *Cureus*. 2024;16(11):e73164.
- [23] Hu R, Niu JY, Wu LN, Sun H, Sun P, Huang JY, et al. Comparison of a tube core and Magill forceps for nasotracheal intubation: A randomised controlled trial. *Trials*. 2021;22(1):697.
- [24] Singh N, Mohanty CR, Sahoo S, Mund M, Raju RB. Comparison of McCoy, Macintosh, and Truview laryngoscope for bougie-aided naso-tracheal intubation: A randomized controlled study. *Int J Crit Illn Inj Sci*. 2024;14(1):03-08.
- [25] Tippmann S, Schäfer J, Winter J, Mühler AK, Schmitz K, Schönfeld, et al. Video versus direct laryngoscopy to improve the success rate of nasotracheal intubations in the neonatal intensive care setting: A randomised controlled trial. *BMJ Paediat Open*. 2023;7(1):e001958.
- [26] Dharanindra M, Jedge PP, Patil VC, Kulkarni SS, Shah J, Iyer S, et al. Endotracheal intubation with king vision video laryngoscope vs macintosh direct laryngoscope in ICU: A comparative evaluation of performance and outcomes. *Indian J Crit Care Med*. 2023;27(2):101-06.
- [27] Araújo B, Rivera A, Martins S, Abreu R, Cassa P, Silva M, et al. Video versus direct laryngoscopy in critically ill patients: An updated systematic review and meta-analysis of randomized controlled trials. *Crit Care*. 2024;28(1):1.
- [28] Sugaya A, Naito K, Goto T, Hagiwara Y, Okamoto H, Watase H, et al. First-pass success of video laryngoscope compared with direct laryngoscope in intubations performed by residents in the emergency department. *Cureus*. 2023;15(10):e47563.
- [29] Sengel N, Karabulut G, Kavuncuoglu D, Selmi NH, Sivgin V, Toprak ME. Nasotracheal intubation performance with the McGrath videolaryngoscope versus macintosh laryngoscope in oral and maxillofacial surgery-An observational study. *Niger J Clin Pract*. 2022;25(5):683-89.
- [30] Kocaturk O, Keles S. Comparison of the HugeMed video laryngoscope with the Macintosh direct laryngoscope for nasotracheal tracheal intubation in children undergoing dental treatment: A randomized controlled clinical study. *Expert Rev Med Devices*. 2024;3:01-08.
- [31] King BJ, Padnos I, Mancuso K, Christensen BJ. Comparing video and direct laryngoscopy for nasotracheal intubation. *Anaesth Prog*. 2020;67(4):193-99.
- [32] Gangishetty A, Jonnavithula N, Geetha S, Muthyala H, Peetha H. A comparison of McGrath video laryngoscope and Macintosh laryngoscope during nasotracheal intubation: A randomised controlled study. *J Perioper. Pract*. 2024;34(12):378-83.
- [33] Rajan S, Chandrasekharan R, Mathew J, Roy RA., Rajkumar R, Paul J. Comparative evaluation of ease of nasotracheal intubation using C-MAC versus Tuoreen Video laryngoscope in adult surgical patients: A randomized study. *J Head Neck Physicia*. 2024;12(1):69-73.
- [34] Singh S, Kumar A, Kumar B, Singh S. Comparison between C-MAC D-blade video laryngoscope and McCoy laryngoscope for nasotracheal intubation in traumatic cervical spine surgery-A randomised controlled trial. *Indian J Anaesth*. 2023;67(9):821-24.
- [35] Merola R, Vargas M, Marra A, Buonanno P, Coviello A, Servillo G, et al. Videolaryngoscopy versus fiberoptic bronchoscopy for awake tracheal intubation: A systematic review and meta-analysis of randomized controlled trials. *J Clin Med*. 2024;13(11):3186.
- [36] Tong JL, Tung A. A randomized trial comparing the effect of fiberoptic selection and guidance versus random selection, blind insertion, and direct laryngoscopy, on the incidence and severity of epistaxis after nasotracheal intubation. *Anaesth Analg*. 2018;127(2):485-89.
- [37] Ozkan AS, Akbas S. Nasotracheal intubation in children for outpatient dental surgery: Is fiberoptic bronchoscopy useful? *Niger J Clin Pract*. 2018;21(2):183-88.
- [38] Tsukamoto M, Kameyama I, Miyajima R, Hitosugi T, Yokoyama T. Alternative Technique for Nasotracheal Intubation Using a Flexible Fiberoptic Scope. *Anaesth Prog*. 2022;69(2):35-37.
- [39] Rhee SH, Yun HJ, Kim J, Karm MH, Ryoo SH, Kim HJ, et al. Risk factors affecting the difficulty of fiberoptic nasotracheal intubation. *J Dent Anaesth Pain Med*. 2020;20(5):293-301.
- [40] Dong Y, Li G, Wu W, Su R, Shao Y. Lightwand-guided nasotracheal intubation in oromaxillofacial surgery patients with anticipated difficult airways: A comparison with blind nasal intubation. *In J Oral Maxillofac Surg*. 2013;42(9):1049-53.
- [41] Fiajoe JE, Stricker P, Gurnaney H, Nishisaki A, Rabinowitz A, Gurwitz A, et al. Ultrasound-guided tracheal intubation a novel intubation technique. *Anesthesiology*. 2012;117(6):1389-91.
- [42] Ahmad I, El-Boghdady K, Bhargava R, Hodzovic I, McNarry AF, Mir F, et al. Difficult Airway Society guidelines for awake tracheal intubation (ATI) in adults. *Anaesthesia*. 2020;75(4):509-28.
- [43] Abdelraziz RA, Ayoub SNB, El-Sherief HM, Shorbagy MS. Pre-emptive epinephrine nebulization prior to nasotracheal intubation for mandibular fracture fixation surgeries: Does it really differ? A randomised controlled clinical trial. *Egypt J Anaesth*. 2023;39(1):348-55.
- [44] Song J. A comparison of the effects of epinephrine and xylometazoline in decreasing nasal bleeding during nasotracheal intubation. *J Dent Anaesth Pain Med*. 2017;17(4):281-87.
- [45] Patel S, Hazarika A, Agrawal P, Jain D, Panda NK. A prospective randomized trial of xylometazoline drops and epinephrine merocele nasal pack for reducing epistaxis during nasotracheal intubation. *J Dent Anaesth Pain Med*. 2020;20(4):223-31.

[46]

Larsen MH, Rosenkrantz O, Creuzburg A, Kristensen MS, Rasmussen LS, Isbye D. Cocaine versus xylometazoline to prevent epistaxis after nasotracheal intubation: A randomized trial. *Acta Anaesthesiol Scand*. 2024;68(9):1192-99.

[47]

Yu HK, Park J, Kang YH, Park HB, Bae SI, Lee SH, et al. Nasal packing with bupivacaine during nasotracheal intubation can reduce intubation-related epistaxis. *Oral Biol Res*. 2021;45(3):107-14.

[48]

Ozkan AS, Akbas S, Toy E, Durmus M. North polar tube reduces the risk of epistaxis during nasotracheal intubation: A prospective, randomized clinical trial. *Curr Ther Res. Clin Exp*. 2018;90:21-26.

[49]

Hoshijima H, Mihara T, Nagumo T, Sato Boku A, Shiga T, Mizuta K. Nasal protection strategy reduces the incidence of nasal pressure injuries during nasotracheal intubation: Meta-analysis with trial sequential analysis. *Medicine*. 2022;101(40):e30638.

[50]

Yang GY, Gao CY, Cai J. Prevention of nasal ala pressure injuries with use of hydroactive dressings in patients with nasotracheal intubation of orthognathic surgery-A randomized controlled trial. *J Wound Ostomy Continence Nurs*. 2020;47(5):484-88.

[51]

Chen J, Chen JL, Yang JN, Chen YX, Liang YF, Lin Y. Investigating the efficacy of hydrocolloid dressing for preventing nasotracheal tube-related pressure injury in the PICU. *Pediatr Crit Care Med*. 2020;21(9):e752-e758.

[52]

Hashimoto M, Sato Boku A, Sento Y, Kamimura Y, Kako E, Okuda M, et al. 3M microfoam™ surgical tape prevents nasal pressure injury associated with nasotracheal Intubation: A randomized double-blind trial. *Medicine*. 2023;102(2):e32679.

[53]

Pedersen J, Schurizek BA, Melsen NC, Juhl B. The effect of nasotracheal intubation on the paranasal sinuses: A prospective study of 434 intensive care patients. *Acta Anaesthesiol Scand*. 1991;35(1):11-13.

[54]

Bach A, Boehrer H, Schmidt H, Geiss HK. Nosocomial sinusitis in ventilated patients: Nasotracheal versus orotracheal intubation. *Anaesthesia*. 1992;47(4):335-39.

[55]

Talesh KT, Gargary RM, Arta SA, Yazdani J, Roshandel M, Ghanizadeh M, et al. Effect of 2% nasal mupirocin ointment on decreasing complications of nasotracheal intubation: A randomized controlled trial. *J Clin Diagn Res*. 2017;11(8):PC08-12.

[56]

Tsukamoto M, Taura S, Kadowaki S, Hitosugi T, Miki Y, Yokoyama T. Risk factors for postoperative sore throat after nasotracheal intubation. *Anaesth Prog*. 2022;69(3):03-08.

[57]

Tachibana N, Niiyama Y, Yamakage M. Less postoperative sore throat after nasotracheal intubation using a fiberoptic bronchoscope than using a Macintosh laryngoscope: A double-blind, randomized, controlled study. *J Clin Anaesth*. 2017;39:113-17.

[58]

Yin F, Zhang TJ. Cardiac arrest by rhino-cardiac reflex during nasotracheal intubation. *Asian J Surg*. 2023;46(6):2358-59.

[59]

Kim H, Lee JM, Lee J, Hwang JY, Chang JE, No HJ, et al. Influence of nasal tip lifting on the incidence of the tracheal tube pathway passing through the nostril during nasotracheal intubation: A randomized controlled trial. *Anaesth Analg*. 2018;127(6):1421-26.

[60]

Won D, Kim H, Chang JE, Lee JM, Min SW, Jung J, et al. Effect of bevel direction on the tracheal tube pathway during nasotracheal intubation: A randomised trial. *Eur J Anaesthesiol*. 2021;38(2):157-63.

PARTICULARS OF CONTRIBUTORS:

1. Doctor, Department of Intensive Care Unit, Huaibei Miners General Hospital, Huaibei, Anhui, China.
2. Doctor, Department of Intensive Care Unit, Huaibei Miners General Hospital, Huaibei, Anhui, China.
3. Doctor, Department of Intensive Care Unit, Huaibei Miners General Hospital, Huaibei, Anhui, China.
4. Professor, Department of Intensive Care Unit, Huaibei Miners General Hospital, Huaibei, Anhui, China.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Yong Wang,
1 Changshan North Road, Huaibei, Anhui, China.
E-mail: wangyongmq@126.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was informed consent obtained from the subjects involved in the study? No
- For any images presented appropriate consent has been obtained from the subjects. No

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Feb 01, 2025
- Manual Googling: May 21, 2025
- iThenticate Software: Jun 02, 2025 (10%)

ETYMOLOGY: Author Origin

EMENDATIONS: 6

Date of Submission: Jan 31, 2025
Date of Peer Review: May 02, 2025
Date of Acceptance: Jun 04, 2025
Date of Publishing: Feb 01, 2026