

A Case of Glutaric Aciduria Type I: An Unusual Cause of Macrocephaly and Developmental Delay

G PREMALAKSHMI¹, SHANTHI RAMESH², JAGDEEP RAMESH³

(CC) BY-NC-ND

ABSTRACT

Glutaric Aciduria Type I (GA-I) is a rare autosomal recessive neurometabolic disorder caused by deficiency of glutaryl-CoA dehydrogenase, leading to accumulation of glutaric acid and 3-hydroxyglutaric acid with subsequent neurotoxicity. A seven-month-old male infant presented with developmental regression, macrocephaly, and recurrent seizures since five months of age. Seizures were characterised by tonic posturing of all four limbs with upward deviation of eyes, followed by postictal drowsiness. Neuroimaging revealed widened Sylvian fissures with a characteristic “bat-wing” appearance, hypoplastic temporal lobes, basal ganglia involvement, and a left-sided subdural haematoma. Tandem mass spectrometry showed elevated glutaryl carnitine (C5-DC: 0.47 $\mu\text{mol/L}$), and genetic analysis identified a homozygous missense variant in the GCDH gene (c.769C>T; p.Arg257Trp), confirming the diagnosis. The child was managed with a lysine-restricted diet, L-carnitine, riboflavin, and antiepileptic therapy. The subdural haematoma was managed conservatively following neurosurgical consultation. This case highlights the importance of early recognition of GA-I in infants presenting with macrocephaly, developmental delay, and characteristic neuroimaging findings, as prompt diagnosis and metabolic therapy can prevent irreversible neurological injury and improve long-term outcomes.

Keywords: Carnitine, Organic acidemia, Seizure, Subdural haematoma, Tandem mass spectrometry

CASE REPORT

A seven-month-old male infant, second born to third-degree consanguineous parents, presented with global developmental delay, regression of milestones, and recurrent seizures since five months of age. The child was apparently well until five months of age, following which he developed multiple seizure episodes. The seizures were characterised by tonic posturing of all four limbs with upward deviation of eyes, lasting less than one minute, followed by postictal drowsiness. He had 3-4 similar episodes during initial hospitalisation and was started on clonazepam. Following the onset of seizures, regression of previously attained milestones was noted. Prior to illness, the child had achieved neck control at three months, rolling over and bimanual grasp at five months, social smile at two months, and recognition of parents at three months. Antenatal and perinatal history was uneventful. The child was born at term by Lower Segment Caesarean Section (LSCS) with a birth weight of 2.5 kg and cried immediately after birth.

Developmental assessment revealed severe global developmental delay with a developmental quotient of approximately 25%. Family history revealed third-degree consanguinity, one healthy sibling, and a history suggestive of sudden infant death in a paternal relative. There was also a history of progressive gait abnormalities in the father and paternal aunt, though no formal evaluation was performed. Genetic evaluation of affected family members could not be performed due to resource limitations. Anthropometry showed weight 7 kg (0 to -2 SD), length 70 cm (+1 to +2 SD), and head circumference 48 cm (+2 to +3 SD), suggestive of macrocephaly.


On examination, the child had dysmorphic facies including depressed nasal bridge and low-set ears [Table/Fig-1]. Neurological examination revealed generalised hypotonia with evolving appendicular hypertonia, brisk deep tendon reflexes, and extensor plantar response appropriate for age. Systemic examination revealed that cardiovascular, respiratory, and abdominal examinations were within normal limits.

The Magnetic Resonance Imaging (MRI) brain showed widened Sylvian fissures with characteristic “bat-wing” appearance, hypoplastic temporal lobes, symmetrical signal changes in bilateral basal ganglia,

and prominent extra-axial spaces [Table/Fig-2]. Computed Tomography (CT) brain revealed a large left convexity subdural haematoma (~16 mm) with midline shift and mass effect [Table/Fig-3]. There was no history of trauma or non accidental injury, and no external signs of injury. Coagulation profile was normal. Despite the presence of a left convexity subdural haematoma (~16 mm) with mild midline shift, the child remained haemodynamically stable without signs of raised



[Table/Fig-1]: Facial dysmorphism showing macrocephaly (arrow) depressed nasal bridge and low-set ears.




along with supportive care. A lysine-restricted diet was advised and continued during follow-up, with avoidance of high-lysine foods such as meat, dairy products, and pulses. After two weeks of hospitalisation, the child was discharged on the same medications. At three-month follow-up, partial seizure control and mild neurological improvement were observed.

At 3-month follow-up, the child continued on the same medications with partial control of seizures. Mild neurological improvement was noted in the form of reduced irritability, improved neck holding, better interaction with caregivers, and decreased frequency of abnormal movements; however, developmental delay persisted.

DISCUSSION

The GA-I is a rare autosomal recessive neurometabolic disorder caused by deficiency of glutaryl-CoA dehydrogenase, resulting in accumulation of glutaric acid and 3-hydroxyglutaric acid. These metabolites exert neurotoxic effects, particularly involving the basal ganglia [2]. Clinically, GA-I commonly presents with macrocephaly, developmental delay, and regression of milestones. Early manifestations may precede acute encephalopathic crises, emphasising the importance of early recognition [3,4]. Several studies have reported cases of GA-I presenting with subdural haematoma, which can mimic non accidental injury. Kölker S et al., described the phenotypic variability of GA-I, including subdural collections in affected children [5]. Boy N et al., further highlighted that early clinical features such as macrocephaly and delayed milestones may be present before neurological deterioration [6]. Previous reports by



intracranial pressure, focal neurological deficits, or rapid neurological deterioration. Following neurosurgical consultation, conservative management with close clinical and radiological monitoring was advised, considering the underlying diagnosis of GAI and the absence of indications for immediate surgical intervention.

Tandem mass spectrometry showed elevated glutaryl carnitine (C5-DC: 0.47 $\mu\text{mol/L}$) [Table/Fig-4]. Genetic analysis identified a homozygous missense variant c.769C>T (p.Arg257Trp) in exon 8 of the GCDH gene, which was classified as likely pathogenic according to the American College of Medical Genetics and Genomics (ACMG) guidelines proposed by Richards S et al., [1]. Metabolic evaluation revealed normal plasma amino acid levels with low free carnitine and elevated glutaryl carnitine levels [Table/Fig-4,5].

Differential diagnoses considered included hypoxic-ischaemic encephalopathy, non accidental injury, mitochondrial disorders, and other organic acidemias, which were excluded based on clinical history, imaging findings, and metabolic evaluation. Hence, a final diagnosis of GA-I with subdural haematoma presenting as developmental regression and seizures was made.

The child was started on Trihexyphenidyl 0.25 mg twice daily, Clonazepam 0.5 mg twice daily, L-carnitine 2.5 mL twice daily, and Riboflavin 100 mg once daily as adjunctive metabolic therapy,

Carnitine	Observed values ($\mu\text{mol/L}$)	Reference range ($\mu\text{mol/L}$)
Free carnitine, C0	6.17 ↓	9.00-69.00
Acetylcarnitine, C2	4.15	2.00-63.00
Propionylcarnitine, C3	0.42	0.08-5.65
Butyrylcarnitine, C4	0.04	0.00-1.30
3-OH-butyrylcarnitine, C4-OH	0.01	0.00-0.50
Isovaleryl/2-methylbutyrylcarnitine, C5	0.07	0.00-0.70
3-OH-isovalerylcarnitine, C5-OH	0.07	0.00-0.80
Glutaryl carnitine, C5-DC	0.47 ↑	0.00-0.38
Tiglylcarnitine, C5:1	0.01	0.00-0.25
Hexanoylcarnitine, C6	0.02	0.00-0.40
Adipyl carnitine, C6DC	0.02	0.00-0.60
Octanoylcarnitine, C8	0.01	0.00-0.48

[Table/Fig-4]: Acylcarnitine profile showing low free carnitine and elevated glutaryl carnitine suggestive of Glutaric Aciduria Type I (GA-I).

Amino acid	Observed values ($\mu\text{mol/L}$)	Reference range ($\mu\text{mol/L}$)
Alanine	231.00	74.00-613.00
Arginine	19.64	0.00-72.00

Argininosuccinic acid	0.06	0.00 - 2.02
Citrulline	9.93	0.00 - 45.00
Glycine	195.95	2.00 - 745.00
Leucine/Isoleucine/Hydroxyproline	63.00	26.00 - 300.00
Methionine	10.73	1.00 - 44.00
Ornithine	77.28	0.00 - 239.00
Phenylalanine	34.69	21.00 - 136.00
Proline	101.80	38.00 - 385.00
Tyrosine	59.90	17.00 - 250.00
Valine	83.21	41.0 - 285.00

[Table/Fig-5]: Plasma amino acid profile showing amino acid levels within normal reference range.

Kölker S et al., Boy N et al., and Harting I et al., have demonstrated that infants with GA-I may present with macrocephaly and subdural haematoma in the absence of trauma, often leading to diagnostic confusion with non accidental injury [5-7].

The pathogenesis of subdural haematoma in GA-I is attributed to stretching of bridging veins due to cerebral atrophy [7]. In such scenarios, absence of trauma, normal coagulation profile, and characteristic neuroimaging findings help differentiate metabolic causes from traumatic aetiologies [8]. Neuroimaging plays a pivotal role in diagnosis. Classical MRI findings include widened Sylvian fissures ("bat-wing" appearance), frontotemporal atrophy, and basal ganglia involvement. These features help distinguish GA-I from hypoxic-ischaemic encephalopathy, mitochondrial disorders, and other organic acidemias [9]. Early diagnosis is crucial, as prompt initiation of therapy can prevent irreversible neurological damage. Management includes a low-lysine diet, L-carnitine supplementation, and riboflavin in selected cases [10,11]. During intercurrent illnesses, emergency protocols with adequate caloric intake and avoidance of catabolism are essential [12]. Studies have shown that early-treated patients have significantly improved neurological outcomes compared to those diagnosed after symptom onset. Regular follow-up with biochemical monitoring is important to ensure treatment adherence and to reduce long-term neurological disability [13,14]. This case underscores the importance of considering metabolic disorders such as GA-I in infants presenting with macrocephaly and subdural haematoma, to avoid misdiagnosis and ensure timely intervention.

CONCLUSION(S)

The GA-1 is a rare but important neurometabolic disorder that should be suspected in infants presenting with macrocephaly, developmental regression, seizures, and subdural haematoma without a history of trauma. Characteristic neuroimaging findings, supported by tandem mass spectrometry and genetic analysis, are essential for establishing

an early diagnosis. Timely initiation of metabolic therapy, including lysine-restricted diet, L-carnitine, and riboflavin supplementation, may significantly reduce neurological deterioration and improve long-term outcomes. This case emphasises the need for increased clinical awareness to prevent delayed diagnosis and misinterpretation as non accidental injury. Early recognition and multidisciplinary management are crucial in minimising irreversible neurological damage and improving quality of life.

REFERENCES

- [1] Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: A joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-24.
- [2] Kölker S, Christensen E, Leonard JV, Greenberg CR, Boneh A, Burlina AB, et al. Diagnosis and management of glutaric aciduria type I - revised recommendations. *J Inher Metab Dis*. 2011;34(3):677-94.
- [3] Boy N, Mengler K, Thimm E, Schiergens KA, Marquardt T, Weinhold N, et al. Newborn screening: A disease-changing intervention for glutaric aciduria type 1. *Ann Neurol*. 2018;83(5):970-79.
- [4] Hoffmann GF, Athanassopoulos S, Burlina AB, Duran M, de Klerk JB, Lehnert W, et al. Clinical course, early diagnosis, treatment, and prevention of disease in glutaryl-CoA dehydrogenase deficiency. *Neuropediatrics*. 1996;27(3):115-23. Doi: 10.1055/s-2007-973761.
- [5] Kölker S, Garbade SF, Boy N, Maier EM, Meissner T, Mühlhausen C, et al. Decline of acute encephalopathic crises in children with glutaryl-CoA dehydrogenase deficiency identified by newborn screening in Germany. *Pediatr Res*. 2007;62(3):357-63. Doi: 10.1203/PDR.0b013e318137a124.
- [6] Boy N, Heringer J, Brackmann R, Bodamer O, Seitz A, Kölker S, et al. Extrastriatal changes in patients with late-onset glutaric aciduria type I highlight the risk of long-term neurotoxicity. *Orphanet J Rare Dis*. 2017;12(1):77. Doi: 10.1186/s13023-017-0612-6.
- [7] Harting I, Neumaier-Probst E, Seitz A, Maier EM, Assmann B, Baric I, et al. Dynamic changes of striatal and extrastriatal abnormalities in glutaric aciduria type I. *Brain*. 2009;132(7):1764-82.
- [8] Strauss KA, Puffenberger EG, Robinson DL, Morton DH. Type I glutaric aciduria, part 1: Natural history of 77 patients. *Am J Med Genet C Semin Med Genet*. 2003;121C(1):38-52.
- [9] Twomey EL, Naughten ER, Donoghue VB, Ryan S. Neuroimaging findings in glutaric aciduria type 1. *Pediatr Radiol*. 2003;33(12):823-30.
- [10] Boy N, Mühlhausen C, Maier EM, Heringer J, Assmann B, Burgard P, et al; Additional individual contributors. Proposed recommendations for diagnosing and managing individuals with glutaric aciduria type I: Second revision. *J Inher Metab Dis*. 2017;40(1):75-101. Doi: 10.1007/s10545-016-9999-9.
- [11] Boy N, Mühlhausen C, Maier EM, Heringer J, Assmann B, Burgard P, et al. Proposed recommendations for diagnosing and managing individuals with glutaric aciduria type I: Second revision. *J Inher Metab Dis*. 2017;40(1):75-101. Doi: 10.1007/s10545-016-9999-9.
- [12] Rainkar M, Kazi ZB, Desai A, Walters C, Rosenberg A, Kishnani PS, et al. High dose IVIG successfully reduces high rhGAA IgG antibody titers in a CRIM-negative infantile Pompe disease patient. *Molecular Genetics and Metabolism*. 2017;122(1-2):76-79. Doi: 10.1016/j.ymgme.2017.05.006.
- [13] Heringer J, Boy N, Ensenaer R, Assmann B, Zschocke J, Harting I, et al. Use of guidelines improves the neurological outcome in glutaric aciduria type I. *Ann Neurol*. 2010;68(5):743-52.
- [14] Kölker S, Boy N, Heringer J, Müller E, Maier EM, Ensenaer R, et al. Complementary dietary treatment using lysine-free, arginine-fortified amino acid supplements in glutaric aciduria type I - A decade of experience. *Mol Genet Metab*. 2012;107(1-2):72-80. Doi: 10.1016/j.ymgme.2012.03.021.

PARTICULARS OF CONTRIBUTORS:

1. Junior Resident, Department of Paediatrics, Sree Balaji Medical College and Hospital, Chennai, Tamil Nadu, India.
2. Professor, Department of Paediatrics, Sree Balaji Medical College and Hospital, Chennai, Tamil Nadu, India.
3. Assistant Professor, Department of Community Medicine, Tagore Medical College and Hospital, Chennai, Tamil Nadu, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Shanthi Ramesh,
Flat No. A2, Ragamalika Apartment, Chennai, Tamil Nadu, India.
E-mail: premagovind21@gmail.com

AUTHOR DECLARATION:

- Financial or Other Competing Interests: None
- Was informed consent obtained from the subjects involved in the study? Yes
- For any images presented appropriate consent has been obtained from the subjects. Yes

PLAGIARISM CHECKING METHODS: [Jain H et al.]

- Plagiarism X-checker: Jan 20, 2026
- Manual Googling: May 28, 2026
- iThenticate Software: May 30, 2026 (5%)

ETYMOLOGY: Author Origin

EMENDATIONS: 7

Date of Submission: Dec 27, 2025

Date of Peer Review: Feb 16, 2026

Date of Acceptance: Jun 01, 2026

Date of Publishing: Jul 01, 2026