

Proportion of Tuberculosis Patients Incurring Catastrophic Health Expenditure in India: A Systematic Review and Meta-analysis Research Protocol

JANMEJAYA SAMAL¹, SHRADDHA BHATIA²

ABSTRACT

Introduction: Globally and at the national level in India, Tuberculosis (TB) programs aim to achieve zero Catastrophic Health Expenditure (CHE) on TB. However, there is evidence through the observational studies reporting CHE incurred by Indian families on TB. Furthermore, the contribution of India to the global TB burden is high both in morbidity and mortality.

Need of the study: A systematic review and meta-analysis will be conducted to estimate the proportion of Indian patients who incur the CHE among Indian TB patients.

Aim: This study aimed to determine the proportion of Indian TB patients incurring CHE, identify factors contributing to CHE among these patients and estimate the mean/median CHE amounts nationally and regionally due to TB in India.

Materials and Methods: Three databases, PubMed, Embase and Scopus, will be used to search for English-language studies reporting the proportion, determinants and mean/median CHE among Indian TB patients. Grey literature will be identified through Google searches and reference screening. Two independent reviewers will screen studies, with disagreements resolved by a senior researcher. Risk of bias will be assessed using the Newcastle-Ottawa Scale (NOS) and the Mixed Methods Appraisal Tool. The review will estimate CHE proportions with 95% confidence intervals and examine predictors, with subgroup, sensitivity and publication bias analyses (I^2 , Egger's, Begg's, trim-and-fill). Results will be presented using forest, funnel, Galbraith and leave-one-out plots. Evidence quality will be assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) tool.

Keywords: High morbidity, Impoverishment, Insurance, Social protection, Social security

INTRODUCTION

The Tuberculosis continues to be a major global public health challenge. Excluding the acute Coronavirus Disease-2019 (COVID-19) peak, TB is the leading cause of death due to a single infectious disease and is listed by World Health Organisation (WHO) as one of the top 10 causes of death around the globe [1]. The burden is disproportionately high in Low and Middle-Income Countries (LMICs). Countries such as India (25%), Indonesia (10%), China (6.5%), the Philippines (6.8%), Pakistan (6.3%), Nigeria (4.8%), Bangladesh (3.6%), South Africa (3.6%) and the Democratic Republic of the Congo (3.9%) made up 71%. Of these, the top five contributed to 55% of the global burden. The report also underlines that India is home to one-third of Multiple Drug Resistant (MDR)/Rifampicin Resistant TB (MDR/RR-TB) cases (32%) and accounts for 28% deaths globally. Approximately, 60% of the difference between the estimated number of individuals worldwide who developed MDR/RR-TB in 2024 (incident cases of MDR/RR-TB) and the number of individuals enrolled in treatment in 2024 was accounted for in five nations.

These nations were India (33%), the Philippines (9.3%), Indonesia (7.3%), China (6.1%) and Pakistan (4.1%) [1].

The WHO's End TB strategy aims to eliminate catastrophic expenditures for households affected by TB, reduce TB incidence by 80% and reduce TB deaths by 90% by 2030 [2]. Health expenditure is considered catastrophic when it jeopardises a household's capacity to cover its basic subsistence needs [3]. Health systems should be designed to minimise out-of-pocket expenses for patients when accessing care. This requires not only expanding universal health coverage to reduce direct medical expenses (such as consultation fees, diagnostic tests and medications), but also strengthening the health system and

improving patient care pathways. A low level of catastrophic expenditure serves as an indicator of a well-functioning health system within an economy [4].

The CHE transpires when a household incurs substantial out-of-pocket healthcare costs, resulting in financial distress or impoverishment [5]. Some defined thresholds of CHE are: 10%, 20% and 40% [6,7]. CHE 10 implies that a household spends more than 10% of its total income on healthcare and so is considered catastrophic at this level [8]. To realise the goal of zero catastrophic expenditure, it is vital to understand the proportion of people still incurring expenditure above a certain threshold and the factors driving it. This will enable us to reshape and manoeuvre the programmatic initiatives.

Most systematic reviews on TB care are either global in scope or focus on selected groups of countries, addressing specific aspects such as TB types or stages, incidence, prevalence, detection strategies and risk of infection [9-15]. Prior to the implementation of the "End TB Strategy," a systematic review of 49 studies conducted across low- and middle-income countries- including India, the Dominican Republic, Bangladesh, Ghana, Vietnam, Thailand, Myanmar, China, Tajikistan and Kenya- reported that the average total cost amounted to 58% of individual annual income and 39% of household income [16]. Another systematic review encompassing 76 studies from low-, middle- and high-income settings found that, despite subsidised TB care, patients continued to incur substantial costs, particularly for hospitalisation and non TB medications. Additionally, patients with drug-resistant TB experienced significantly higher financial burden [17]. A review of 29 studies predominantly based in India, China, Indonesia, Uganda and South Africa found that approximately half of households (43%) affected by TB experienced CHE when applying

a 10% threshold of income [18]. With a higher threshold of 20%, 40% of the 196 pulmonary TB patients surveyed in Uganda faced catastrophic costs [19]. Thus, the financial burden associated with seeking TB diagnosis and treatment remains a substantial barrier to care at the global level.

Given the significant share of India in TB cases and related deaths, it is imperative to delve into the catastrophic cost scenario. Despite the provision of free TB care through the National TB Elimination Program (NTEP), over one-third of patients in Puducherry experienced catastrophic expenses [20]. Among the rural patients, approximately half consult private clinics and 20% purchase anti-TB drugs from private pharmacies, which increase the out-of-pocket expenditure [21]. A 30-60% of the patients with drug-susceptible TB across Assam, Maharashtra, Tamil Nadu and West Bengal incurred catastrophic expenditure [22]. The TB disproportionately affects people in the working age group in India, which further increases the income and productivity loss [23,24]. Addressing TB in India is essential for global TB elimination efforts, as progress here will significantly influence global outcomes.

Despite the availability of free diagnosis and treatment under the National TB Elimination Program, TB continues to impose a significant financial burden on Indian families. Many Indian TB patients lose livelihood support, direct and indirect costs related to diagnosis and treatment, which pushes them into poverty. Given this understanding, estimating the CHE incurred by Indian TB patients has become critical for monitoring both the NTEP target and WHO's End-TB targets. Nonetheless, the evidence on CHE remains fragmented and varies across different settings in India. Although several observational studies are available [21,24-27], there is no comprehensive systematic review or meta-analysis that estimates the proportion of Indian patients experiencing CHE, identifies its key drivers, or quantifies the mean/median CHE attributable to TB in India. The present systematic review will fill the gap by synthesising existing data and a robust national estimate to inform policies and social protection schemes.

Primary objectives: To determine the pooled proportion of Indian patients incurring CHE due to TB.

Secondary objectives: To identify the factors contributing to the CHE among Indian TB patients and to estimate the mean/median total CHE (regional/national) due to TB in India.

MATERIALS AND METHODS

According to Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P), this protocol was registered with PROSPERO on 1st December 2025 (ID: CRD420251242934). The PRISMA-P checklist was used to write this protocol [28]. The Population, Intervention/Exposure, Comparison, Outcome and Study Designs (PICOS) framework will be used to finalise which studies to include and which to exclude. The search process preceded writing this protocol.

The literature search will be conducted across three databases: PubMed, Embase and Scopus [Table/Fig-1].

Inclusion and Exclusion criteria:

Population: The analysis will include studies conducted among all types of TB (pulmonary TB, extra-pulmonary TB, drug-sensitive TB and drug-resistant TB) in Indian settings, encompassing populations of all ages, sexes and socio-economic groups.

Exclusion criteria will involve excluding studies that were conducted outside India and populations other than TB patients.

Intervention/Exposure: Studies will be included that report the experience of CHE (both direct and indirect costs and/or a combination of both) among Indian TB patients and will consider the CHE at any threshold level, such as 10%, 20%, or 40%.

Name of the database	Search details	Total articles
PubMed	{("Tuberculosis (TB)"(MeSH Terms) OR "Tuberculosis (TB)" (All Fields) OR "TB"(All Fields)) AND ("Catastrophic Health Expenditure (CHE)" (All Fields) OR "catastrophic health spending"(All Fields) OR "catastrophic cost"(All Fields) OR "out-of-pocket expenditure"(All Fields) OR "financial burden"(All Fields) OR "health expenditure"(All Fields) OR "healthcare expenditure"(All Fields) OR "economic burden"(All Fields) OR "cost of illness"(All Fields)) AND ("India"(MeSH Terms) OR "India"(All Fields) OR "Indian"(All Fields))}	190
Scopus	TITLE-ABS-KEY (("TB" OR "TB") AND ("Catastrophic Health Expenditure (CHE)" OR "catastrophic health spending" OR "catastrophic cost" OR "out-of-pocket expenditure" OR "financial burden" OR "health expenditure" OR "healthcare expenditure" OR "economic burden" OR "cost of illness")) AND ("India" OR "Indian"))	187
Embase	('tuberculosis'/exp OR tuberculosis:ti,ab OR TB:ti,ab) AND ('Catastrophic Health Expenditure (CHE) ':ti,ab OR 'catastrophic health spending':ti,ab OR 'catastrophic cost':ti,ab OR 'out of pocket expenditure':ti,ab OR 'financial burden':ti,ab OR 'health expenditure':ti,ab OR 'healthcare expenditure':ti,ab OR 'economic burden':ti,ab OR 'cost of illness':ti,ab) AND ('india'/exp OR india:ti,ab OR indian:ti,ab)	117
Total		494

[Table/Fig-1]: Search strategy for systematic literature review and meta-analysis.

Comparator: Since the objective of the study is related to the proportion of Indian patients incurring the CHE due to TB, the comparator is not applicable for the present study.

Outcomes: Studies will be included that report the proportion of Indian TB patients incurring CHE, drivers of CHE in Indian TB patients and mean/median expenditure due to TB.

Study design: Study designs include hospital- and community-based observational studies (prospective, retrospective, cross-sectional, case-control and cohort), as well as qualitative and interventional studies.

Study Procedure

Data selection, extraction and management: Article selection will take place in two stages. The initial screening at the title and abstract stage will be completed by two reviewers and then the disagreements will be mutually discussed. Full texts will be obtained for all articles reviewed. The second stage will consist of full-text screening.

The reviewers will independently divide and screen the studies. During the full-text screening stage, reasons for exclusion will be documented. Any disagreements will be resolved through discussion and if consensus is not reached, a third reviewer will adjudicate. In cases where the same study is reported in multiple publications, these will be linked and recorded as a single study.

Data will be extracted using a structured form developed in line with the study objectives and research questions. The form will undergo a pilot testing and calibration exercise before use. Subsequently, each reviewer will independently extract data using the standardised form. If additional information is required, attempts will be made to contact the original study authors, with a maximum of two follow-up attempts.

Extracted data will include, at a minimum, the following: study and reviewer identifiers; study characteristics (author, year of publication, study design); setting characteristics (national or sub-national—state, district and so on); population characteristics (adult or juvenile, age, sex, type of TB, type of facility sought, socio-economic status and other relevant information); outcomes (proportion of sample facing CHE); mean/median expenditure and drivers of catastrophic expenditure if reported; other study specifications (co-morbidities); and general comments.

Risk of bias: The risk of bias in the included studies will be evaluated using the NOS and the Modified-NOS for cohort and case-control studies, respectively, as well as for cross-sectional studies [29]. The Mixed Methods Appraisal Tool will be utilised for mixed-method investigations [30]. Two reviewers will check for bias and disagreements will be settled mutually. The scoring scheme for NOS is: 7-9 stars: low, 5-6 stars: moderate and less than 5 stars: high-risk of bias. The modified NOS will be classified into 6-7 stars, 4-5 stars and less than 4 stars, showing low, moderate and high-risk of bias, respectively. Since NOS does not cover all types of bias, such as reporting bias and selective outcome reporting the limitation section will acknowledge this [29]. The certainty of evidence will be assessed using the Grading of Recommendations Assessment, Development and Evaluation (GRADE)-aligned approach for the cross-sectional studies [31]. Analysis will be conducted using Stata-18 (Stata Corp LLC, College Station, TX, USA).

Outcome Measures

The main outcomes for which data will be sought are as follows.

- **Proportion of Indian TB patients who have incurred the CHE due to TB:** Proportion of TB patients incurring catastrophic cost, where cost includes direct medical cost, such as medicines, consultation, indirect cost, such as transportation, food, accommodation and loss of wages.
- **Drivers of CHE among TB patients in India:** Factors leading to CHE, such as hospitalisation and private consultation.
- **Mean/Median CHE due to TB in India:** Mean or Median total cost (includes direct and indirect).

STATISTICAL ANALYSIS

The pooled proportion of Indian TB patients facing CHE will be calculated using a random-effects model with a 95% confidence interval, considering heterogeneity. A fixed-effects model will be applied if $I^2 < 25\%$; if $I^2 > 75\%$, the random-effects model will be used. For I^2 values between 25%-75%, sample size and study design will also be assessed. Graphical representations will include a forest plot, funnel diagram, Galbraith plot, leave-one-out meta-analysis and a bubble plot for various analyses. Publication bias will be evaluated using Egger's and Begg's tests, with the meta-trim-and-fill method applied if more than 10 studies are included. Model fit and residual heterogeneity will be evaluated using the Q statistic and τ^2 values, with statistical significance set at $p < 0.05$.

Acknowledgement

The authors gratefully acknowledge the financial support of SRM School of Public Health, Faculty of Medicine and Health Sciences, SRMIST, Kattankulathur for bearing the defrayed costs of publishing this article.

REFERENCES

- [1] WHO. Global Tuberculosis Report 2025 [Internet]. 2025 [cited 2025 Dec 8]. Available from: <https://www.who.int/teams/global-programme-on-tuberculosis-and-lung-health/tb-reports/global-tuberculosis-report-2025>.
- [2] The End TB Strategy [Internet]. [cited 2026 Mar 17]. Available from: <https://www.who.int/teams/global-programme-on-tuberculosis-and-lung-health/the-end-tb-strategy>.
- [3] Squire S, Thomson R, Namakhoma I, El Sony A, Kritski A, Madan J. Catastrophic care-seeking costs as an indicator for lung health. *BMC Proc*. 2015;9(10):S4. doi:10.1186/1753-6561-9-S10-S4.
- [4] Reddy US. Measurement of catastrophic health expenditure in india: A systematic review and meta-analysis. *Appl Health Econ Health Policy*. 2024;22(4):471-83. Doi: 10.1007/s40258-024-00885-1. Epub 2024 May 10. PubMed PMID: 38727917.
- [5] Xu K, Evans DB, Kawabata K, Zeramdlini R, Klavus J, Murray CJ. Household catastrophic health expenditure: A multicountry analysis. *The Lancet*. 2003;362(9378):111-17. Doi: 10.1016/S0140-6736(03)13861-5 PubMed PMID: 12867110.
- [6] Xu K. Distribution of health payments and catastrophic expenditures Methodology [Internet]. WHO; 2005 [cited 2026 Feb 22]. Available from: <https://www.who.int/publications/i/item/EIP-FER-DP.05.2>.
- [7] van Doorslaer E, O'Donnell O, Rannan-Eliya RP, Somanathan A, Adhikari SR, Garg CC, et al. Catastrophic payments for health care in Asia. *Health Econ*. 2007;16(11):1159-84. Doi: 10.1002/hec.1209 PubMed PMID: 17311356.
- [8] Ranson MK. Reduction of catastrophic health care expenditures by a community-based health insurance scheme in Gujarat, India: Current experiences and challenges. *Bull World Health Organ*. 2002;80(8):613-21. PubMed PMID: 12219151; PubMed Central PMCID: PMC2567585.
- [9] Beijer U, Wolf A, Fazel S. Prevalence of tuberculosis, hepatitis C virus, and HIV in homeless people: A systematic review and meta-analysis. *Lancet Infect Dis*. 2012;12(11):859-70. Doi: 10.1016/S1473-3099(12)70177-9 PubMed PMID: 23265606; PubMed Central PMCID: PMC3494003.
- [10] Gao J, Zheng P, Fu H. Prevalence of TB/HIV co-infection in countries except China: A systematic review and meta-analysis | PLOS One [Internet]. 2013 [cited 2025 Dec 5]. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0064915>.
- [11] Uden L, Barber E, Ford N, Cooke GS. Risk of tuberculosis infection and disease for health care workers: An updated meta-analysis. *Open Forum Infect Dis*. 2017;4(3):ofx137. Doi: 10.1093/ofid/ofx137 PubMed PMID: 28875155; PubMed Central PMCID: PMC5575844.
- [12] Noykhovich E, Mookherji S, Roess A. The risk of tuberculosis among populations living in slum settings: A systematic review and meta-analysis. *J Urban Health*. 2019;96(2):262-75. Doi: 10.1007/s11524-018-0319-6 PubMed PMID: 30341562; PubMed Central PMCID: PMC6458189.
- [13] Kuupiel D, Vezi P, Bawontuo V, Osei E, Mashamba-Thompson TP. Tuberculosis active case-finding interventions and approaches for prisoners in sub-Saharan Africa: A systematic scoping review. *BMC Infect Dis*. 2020;20(1):570. Doi: 10.1186/s12879-020-05283-1 PubMed PMID: 32758165; PubMed Central PMCID: PMC7405346.
- [14] Cords O, Martinez L, Warren JL, O'Marr JM, Walter KS, Cohen T, et al. Incidence and prevalence of tuberculosis in incarcerated populations: A systematic review and meta-analysis. *Lancet Public Health*. 2021;6(5):e300-e308. Doi: 10.1016/S2468-2667(21)00025-6 PubMed PMID: 33765455; PubMed Central PMCID: PMC8168455.
- [15] Litvinenko S, Magwood O, Wu S, Wei X. Burden of tuberculosis among vulnerable populations worldwide: An overview of systematic reviews. *The Lancet Infectious Diseases*. 2023;23(12):1395-407. Doi: 10.1016/S1473-3099(23)00372-9.
- [16] Tanimura T, Jaramillo E, Weil D, Raviglione M, Lönnroth K. Financial burden for tuberculosis patients in low- and middle-income countries: A systematic review. *European Respiratory Journal*. 2014;43(6):1763-75. Doi: 10.1183/09031936.00193413 PubMed PMID: 24525439.
- [17] D'Silva OA, Lancione S, Ananthakrishnan O, Addae A, Shrestha S, Alsdurf H, et al. The catastrophic cost of TB care: Understanding costs incurred by individuals undergoing TB care in low-, middle-, and high-income settings – A systematic review. *PLOS Global Public Health*. 2025;5(4):e0004283. Doi: 10.1371/journal.pgph.0004283.
- [18] Ghazy RM, El Saeh HM, Abdulaziz S, Hammouda EA, Elzorkany AM, Khidr H, et al. A systematic review and meta-analysis of the catastrophic costs incurred by tuberculosis patients. *Sci Rep*. 2022;12(1):558. Doi: 10.1038/s41598-021-04345-x.
- [19] Walcott RL, Ingels JB, Corso PS, Zalwango S, Whalen CC, Sekandi JN. There's no such thing as a free TB diagnosis: Catastrophic TB costs in Urban Uganda. *Global Public Health*. 2020;15(6):877-88. Doi: 10.1080/17441692.2020.1724313 PubMed PMID: 32027555.
- [20] Prasanna T, Jeyashree K, Chinnakali P, Bahurupi Y, Vasudevan K, Das M. Catastrophic costs of tuberculosis care: A mixed methods study from Puducherry, India. *Global Health Action*. 2018;11(1):1477493. Doi: 10.1080/16549716.2018.1477493 PubMed PMID: 29902134.
- [21] Panda A, Behera BK, Mishra A. Financial hardship of tuberculosis patients registered under National Tuberculosis Elimination Programme (NTEP) in rural India: A longitudinal study. *Indian J Tuberc*. 2024;71 Suppl 2:S229-S236. Doi: 10.1016/j.ijtb.2024.01.007.
- [22] Chatterjee S, Das P, Stallworthy G, Bhambure G, Munje R, Vassall A. Catastrophic costs for tuberculosis patients in India: Impact of methodological choices. *PLOS Glob Public Health*. 2024;4(4):e0003078. Doi: 10.1371/journal.pgph.0003078 PubMed PMID: 38669225; PubMed Central PMCID: PMC11051603.
- [23] Malik TZ, Ahmed F, Roy S, Agarwalla R, Pathak R. Catastrophic costs of Tuberculosis in patients registered under the National Tuberculosis Elimination Programme in South-East Delhi, India. *Clinical Epidemiology and Global Health*. 2025;34. Doi: 10.1016/j.cegh.2025.102075.
- [24] Yadav J, John D, Allarakha S, Menon GR. Rising healthcare expenditure on tuberculosis: Can India achieve the End TB goal. *Trop Med Int Health*. 2021;26(11):1256-75. Doi: 10.1111/tmi.13648 PubMed PMID: 34192385.
- [25] Sarin R, Vohra V, Singla N, Thomas B, Krishnan R, Muniyandi M. Identifying costs contributing to catastrophic expenditure among TB patients registered under RNTCP in Delhi metro city in India. *Indian J Tuberc*. 2019;66(1):150-57. Doi: 10.1016/j.ijtb.2018.10.009 PubMed PMID: 30797274.
- [26] Rupani MP, Vyas S, Shah IA. Cohort study on association between catastrophic costs and unfavorable tuberculosis treatment outcomes among TB-HIV and TB-diabetes comorbid patients in India. *BMC Public Health*. 2024;24(1):2028. Doi: 10.1186/s12889-024-19609-0 PubMed PMID: 39075416; PubMed Central PMCID: PMC11285260.
- [27] Dutta A, Mandal S, Taraphdar P, Das S. Catastrophic cost and coping strategies of tuberculosis patients in a block of Purba Bardhaman District, West Bengal: A cross-sectional study. *Indian J Tuberc*. 2025;72(3):354-57.

[28]

Shamseer L, Moher D, Clarke M, Gherzi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: Elaboration and explanation. *BMJ*. 2015;350:g7647. Doi: 10.1136/bmj.g7647 PubMed PMID: 25555855.

[29]

Carra MC, Romandini P, Romandini M. Risk of bias evaluation of cross-sectional studies: Adaptation of the Newcastle-Ottawa Scale. *J Periodontal Res*. 2025 Apr 28. Doi: 10.1111/jre.13405 PubMed PMID: 40293188.

[30]

Hong QN, Fàbregues S, Bartlett G, Boardman F, Cargo M, Dagenais P, et al. The Mixed Methods Appraisal Tool (MMAT) version 2018 for information professionals and researchers. *Education for Information*. 2018;34(4):285-91. Doi: 10.3233/EFI-180221.

[31]

Prasad M. Introduction to the GRADE tool for rating certainty in evidence and recommendations. *Clinical Epidemiology and Global Health*. 2024;25:101484. Doi: 10.1016/j.cegh.2023.101484.

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, School of Public Health, SRM Institute of Science and Technology (SRMIST), Kattankulathur, Chennai, Tamil Nadu, India.

2. Research Scholar, Department of Economics, PPN (PG) College, CSJM University, Kanpur, Uttar Pradesh, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Shraddha Bhatia,
7/212, Ground Floor, Sai Leela Apartment, Swaroop Nagar,
Kanpur-208002, Uttar Pradesh, India.
E-mail: bhatia10893@gmail.com

PLAGIARISM CHECKING METHODS: [\[Jain H et al.\]](#)

• Plagiarism X-checker: Jan 05, 2026

• Manual Googling: Apr 28, 2026

• iThenticate Software: Apr 30, 2026 (9%)

ETYMOLOGY: Author Origin

EMENDATIONS: 6

AUTHOR DECLARATION:

• Financial or Other Competing Interests: None

• Was Ethics Committee Approval obtained for this study? No

• Was informed consent obtained from the subjects involved in the study? No

• For any images presented appropriate consent has been obtained from the subjects. NA

Date of Submission: [Jan 04, 2026](#)

Date of Peer Review: [Feb 10, 2026](#)

Date of Acceptance: [May 02, 2026](#)

Date of Publishing: [Sep 01, 2026](#)

4

Journal of Clinical and Diagnostic Research. 2026 Sep, Vol-20(9): LK01-LK04