

Antiulcer Activity of Ethanolic Extract of *Salvadora indica* (W.) Leaves on Albino Rats

SAROJ KUMAR SAHOO¹, HIMANSHU BHUSAN SAHOO², D. PRIYADARSHINI³, G. SOUNDARYA⁴, CH. KISHORE KUMAR⁵, K. USHA RANI⁶

ABSTRACT

Background: Ulcer can be developed inside the inner lining of the stomach (gastric ulcer) or the small intestine (duodenal ulcer). Both the ulcers are also cumulatively referred as peptic ulcers. It affects nearly 10% of world population.

Aim: To investigate the antiulcer activity of ethanolic extract of *Salvadora indica* W. leaves (ESIL) on albino rats.

Materials and Methods: The present study was carried by pylorus ligation, ethanol and cysteamine induced ulcer models in albino rats. The antiulcer activity of ESIL (150, 300 and 600 mg/kg p.o. for 7 days) was compared with standard drugs (Ranitidine). In pyloric ligation induced ulcer model, the studied parameters were gastric volume, pH, total acidity, free acidity, and ulcer index whereas in ethanol and cysteamine induced ulcer

model, the ulcer index was determined for severity of ulcers. The parameters studied were ulcer index, gastric juice volume, pH, free acidity and total acidity.

Results: In pyloric ligation model; the volume of gastric content, total/free acidity and pepsin activity was significantly decreased at $p < 0.05$ and $p < 0.01$ and pH of the gastric juice was significantly increased at $p < 0.05$ and $p < 0.01$ in ESIL treated groups as compared to control group. All the doses of ESIL showed dose dependent antiulcer effect as well as significant ($p < 0.05$ and $p < 0.01$) reduction in the ulcer index as compared to control group in all the experimental models.

Conclusion: The results of the study indicate that the ESIL have better potential against ulcer which supports the traditional claims in folklore medicine.

Keywords: Antiulcer effect, Cysteamine, Ethanol, Pylorus ligation, Ulcer index

INTRODUCTION

Ulcer is a major disease of gastrointestinal system which affects 10% of the world population with different aetiologies. Chronic alcohol intake, smoking, excessive stress, chronic usage of non-steroidal anti-inflammatory drugs and *H. pylori* bacterial infection are the crucial causes of peptic ulcer characterized by inflammation, mucosal bleeding and abdominal pain in patients [1,2]. These ulcers can develop when the imbalance occurs between the gastroprotectives (mucus, bicarbonate and prostaglandins) and aggressives (acid, pepsin, bile salts and *Helicobacter pylori* bacteria) [3]. The recent approach to peptic ulcer is managed by inhibition of gastric acid secretion, promotion of gastro-protection, blocking apoptosis and stimulation of epithelial cell proliferation for effective healing. The conventional drugs used in the treatment of ulcer include histamine receptor antagonists, prostaglandins analogues, proton pump inhibitors, cytoprotective agents, antacids and anticholinergics, but most of these drugs produce undesirable side effects or drug interactions and may even alter biochemical mechanisms of the body upon chronic usage. Hence, herbal medicines are generally used in such chronic cases, wherein drugs are required to be used for long periods [4].

Salvadora indica W. is commonly known as 'jhak' in Hindi, belonging to Salvadoraceae family. It is native from Africa, Saudi Arabia, India, Srilanka, Pakistan, Sudan, Egypt and Ceylon [5]. The plant was famous due to its various biological properties as abrasives, astringents and antiseptics. Traditionally, it was used as an important ingredient for tooth care product by Arabian people [6]. The stem of the plant was scientifically proved as antihyperlipidemic and antitumor potential on rodents [7]. The leaves and roots of *S. indica* had been used for the treatment of peptic ulcers as well as syphilitic or other inflammatory ulcers by Ayurvedic practitioners [7-9]. However, there is no scientific evidence on prevention of peptic ulcers of *S. indica* against above traditional claims so far.

Hence, the present study was conducted with the aim to investigate the antiulcer activity of Ethanolic Extract of *Salvadora Indica* W. leaves (ESIL) in albino rats.

MATERIALS AND METHODS

In-vivo antiulcer study was performed from October 2015 to November 2015 and the ethical approval was permitted from Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.

Drugs and Chemicals

Ethanol (Bengal Chemicals, Kolkatta, India), Cysteamine hydrochloride (Hi-media, Mumbai, India) and Ranitidine (Kopran Pharma Ltd., Mumbai, India) were obtained. All other chemicals and solvents used in this study were of analytical grade.

Plant material

The fresh leaves of *S. indica* were collected in the month of July 2015 from the local areas of Srikakulam, Andhra Pradesh, India. The plant was identified and authenticated by Dr. K. M. Chetty, Sri Venkateswara University, Tirupathi, where the voucher specimen was deposited for further reference.

Preparation of Plant Extract

The leaves of *S. indica* were washed thoroughly in water to remove foreign matter and allowed to shade dry with a relative humidity of 40–45%. Then, the leaves were powdered in roller grinder and passed through a sieve (No. 40). Then, the fine powder (Approx. 150 gm) was defatted with petroleum ether and extracted with 1 litre of 95% ethanol at room temperature by using Soxhlet apparatus for 72 hour. The resultant extract was filtered and concentrated in a rotary evaporator under reduced pressure to obtain a thick semi solid brown paste, which was stored at -20°C until required. The yield of the extract was found to be 11.87 %w/w.

Phytochemical Screening

The preliminary phytochemical screening was performed with the ethanolic extract of *S. indica* leaves (ESIL) for the detection of various phytochemicals [10].

Experimental Animals

Adult albino rats (40 numbers) of 130-160g of body weight of either sex were procured from the animal house of Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India. The study protocol was approved from the Institutional Animal Ethics Committee (IAEC) under the reference no. 1427/PO/a/11/CPCSEA and CPCSEA guidelines were adhered during the maintenance and experiment. All animals were maintained under standard husbandry conditions with food and water ad libitum.

Acute Oral Toxicity Study

Healthy adult male albino rats were fasted overnight prior to the experiment. Different doses (50-2000 mg/kg, P.O) of the ethanolic extract of *S. indica* were administered to each group of rats (Each group carries 8 rats) and they were observed continuously for 1 hour and then at half-hourly intervals for 4 hour, for any gross behavioural changes and further up to 72 hour, followed 14 days for any mortality as per the OECD (Organization for Economic Co-operation and Development) Guideline 425 [11]. The leaf extract of *S. indica* was found to be non-toxic up to the maximum dose of 2000 mg/kg body weight. Dose selected for antiulcer evaluation was 150, 300 and 600 mg/kg respectively [11].

Animal Grouping and Treatment Schedule for Antiulcer Study

In all the experimental models, male albino rats were selected and divided into five groups of six animals each. Animals were fasted for 24 hour before the study, but had free access to water. Group I treated as vehicle control, received only distilled water; group II, III and IV treated as treatment groups, received the graded dose of ethanol extract of *S. indica* (ESIL) at 150, 300 and 600 mg/kg, (P.O.) for 7 days (once in a day) respectively and group V as standard group, received ranitidine 50 mg/kg (P.O.) [Table/Fig-1].

Pyloric Ligation Induced Gastric Ulceration

In this method, albino rats were fasted in individual cages for 24 hour ESIL, reference drug and control vehicle was administered 1 hour prior to pyloric ligation. Then the pre-treated animals were anaesthetised by anaesthetic ether; the abdomen was opened by a small midline incision below the xiphoid process. The pyloric portion of the stomach was ligated without causing any damage to its blood vessels. The stomach was isolated carefully and the abdominal wall was sealed by interrupted sutures. The animals were deprived of water during the postoperative period. Four hours after ligation, the stomach was dissected out and contents were collected into clean tubes. The volume, pH and total acid content of gastric juice were determined. The contents were centrifuged, filtered and subjected to titration for estimation of total acidity. From the supernatant, aliquots (1 ml each) were taken for the determination of pH, total or free acidity and pepsin activity [12]. Each stomach was examined for lesions in the fore stomach portion and indexed according to severity [13].

The numbers of ulcers were counted and scoring of ulcer was made as follows: Normal colored stomach (0), Red coloration (0.5), Spot ulcer (1), Haemorrhagic streak (1.5), Deep ulcers (2) and Perforation (3). Mean ulcer score for each animal was expressed as ulcer index [14]. Ulcer index (U_i) was measured by using following formula: $U_i = U_N + U_S + U_p \times 10^{-1}$ Where, U_i (Ulcer Index); U_N (Average number of ulcers per animal); U_S (Average number of severity score); U_p (Percentage of animals with ulcers). The percentage inhibition of ulceration was calculated and compared with control.

Ethanol Induced Mucosal Damage in Rats

The rats were fasted for 24 hours before the experiment. After 1 hour of administration of ESIL, ranitidine and vehicle control treatment, 1ml of absolute ethanol (0.5 ml/100g) was orally administered to each rat of every group. After 1 hour, the animals were sacrificed with excess of anaesthetic ether and stomach was opened along the greater curvature, cleared of residual matter with saline and the inner surface was examined for severity of ulceration [15]. Ulcer index and % ulcer protection were calculated.

Cysteamine Induced Duodenal Ulceration

Cysteamine HCl (400 mg/kg, P.O. in 10% aqueous solution) was administered twice in a day at an interval of 4 hour to produce duodenal ulcers in rats. The *S. indica* or reference drug or control vehicle was administered 30 min before each dose of cysteamine HCl. After 20 hours of last dose of cysteamine, the animals were sacrificed and the duodenum was isolated carefully and opened the portion of the antimesenteric side. The ulcer score was obtained by measuring the dimensions of the duodenal ulcer(s) in square millimeters and ulcer index was determined using the method described earlier [12].

S. N.	Groups	Sample Size
1	Vehicle control (with water)	6
2	Ethanolic extract of <i>S. indica</i> (ESIL) at 150 mg/kg,	6
3	Ethanolic extract of <i>S. indica</i> (ESIL) at 300 mg/kg	6
4	Ethanolic extract of <i>S. indica</i> (ESIL) at 600 mg/kg	6
5	Standard group ranitidine 50 mg/kg (p.o.).	6

[Table/Fig-1]: Groups of experimental animals for antiulcer study.

STATISTICAL ANALYSIS

The results were expressed as the Mean \pm SD for each group. Statistical differences were evaluated using a one-way analysis of variance (ANOVA) followed by Dunnet's t-test. Results were considered to be statistically significant at $p < 0.05$.

RESULTS

Preliminary Phytochemical Screening

The ESIL showed the presence of flavonoids, alkaloids, phenolics, terpenoids, tannins, saponins, carbohydrates and amino acids.

Effect of the ESIL on Pylorus Ligation Induced Gastric Ulceration

In pylorus ligated rats, the volume of gastric content, pH, pepsin activity, total and free acidity are shown in [Table/Fig-2]. In ESIL treated groups, the volume of acid secretion, total acidity and pepsin activity was decreased and pH of the gastric juice was increased compared to ulcer control group. The effects of ethanolic extract of *S. indica* on acid parameters showed significant ($p < 0.01$ and $p < 0.05$) effect at 150, 300 and 600 mg/kg doses compared to ulcer control animals. In addition, *S. indica* showed a dose dependent curative ratio compared to ulcer control groups. The extracts exhibited an inhibition percentage of 37.65, 45.88 and 67.48 at doses of 150, 300 and 600 mg/kg doses respectively; and, the standard drug, ranitidine, exhibited an inhibition percentage of 75.10. ESIL at 600 mg/kg was shown equipotent antiulcer activity with ranitidine.

Effect of the ESIL on Ethanol Induced Gastric Ulceration

The animals pre-treated with ESIL groups showed significant ($p < 0.01$) as well as dose dependent inhibition of ulcer index as compared to control group [Table/Fig-3]. ESIL showed 41.76, 50.06 and 61.30 % ulceration inhibition at 150, 300 and 600 mg/kg respectively whereas ranitidine showed 69.47% ulceration inhibition [Table/Fig-4].

Treatment	Gastric Content (ml)	pH	Acidity (mEq/lit)		Pepsin activity (Per ml/h)
			Total	Free	
Vehicle Control	10.6 ± 0.34	3.41 ± 0.12	121.3 ± 1.30	97.3 ± 0.52	3.75 ± 0.15
ESIL 150 mg/kg	4.26 ± 0.62*	4.80 ± 0.11*	71.16 ± 1.42*	64.39 ± 1.27	3.42 ± 0.90*
ESIL 300 mg/kg	4.86 ± 0.20*	5.08 ± 0.15*	63.13 ± 1.60*	57.7 ± 0.16*	3.17 ± 0.22*
ESIL 600 mg/kg	5.53 ± 0.15**	5.62 ± 0.08**	54.17 ± 0.54**	47.5 ± 0.05*	2.93 ± 0.25*
Ranit. 150 mg/kg	5.68 ± 0.18**	5.86 ± 0.18**	45.2 ± 0.54**	39.2 ± 0.04**	2.84 ± 0.77**


[Table/Fig-2]: Effect of ESIL on gastric parameters by pyloric ligation induced ulceration in rats.

Values are expressed as Mean ± SEM, n= 6, *p < 0.05 and ** p < 0.01 when compared with vehicle control group. (Statistically analysed by one-way ANOVA followed by Dunnet's t-test).

Treatment	Ulcer index		
	Pyloric ligation	Ethanol	Cysteamine
Vehicle Control	4.86 ± 0.21	7.83 ± 0.46	4.32 ± 0.14
ESIL 150 mg/kg	3.03 ± 0.54*	4.56 ± 1.08*	2.88 ± 0.98*
ESIL 300 mg/kg	2.63 ± 0.74*	3.91 ± 0.15*	2.53 ± 0.42*
ESIL 600 mg/kg	1.58 ± 0.63**	3.03 ± 0.36**	1.49 ± 0.87**
Ranitidine 150mg/kg	1.21 ± 0.43**	2.39 ± 0.08**	1.38 ± 0.21**

[Table/Fig-3]: Effect of ESIL on ulcer index by pyloric ligation, ethanol and cysteamine induced ulcers in rats.

Values are expressed as Mean ± SEM, n= 6, *p < 0.05 and ** p < 0.01 when compared with vehicle control group. (Statistically analysed by one-way ANOVA followed by Dunnet's t-test).



[Table/Fig-4]: Effect of ESIL on percentage of ulcer inhibition by pyloric ligation, ethanol and cysteamine induced ulcers in rats.

Values are expressed as Mean ± SEM, n= 6, *p < 0.05 and ** p < 0.01 when compared with control group. (Statistically analysed by one-way ANOVA followed by Dunnet's t-test).

Effect of ESIL on Cysteamine Induced Duodenal Ulcers

In the cysteamine induced duodenal ulcers, oral administration of ESIL showed a reduction in ulcer index in a dose dependent manner. The percentage ulcer inhibition was 33.33, 41.43 and 65.5 for 150, 300 and 600 mg/kg respectively and that of the standard was found to be 68.05 [Table/Fig-3], [Table/Fig-4].

DISCUSSION

The cause of gastric ulcer is due to stress induced increase in gastric acid (HCl) secretion and these acid secretions promote ulceration due to exposure of the unprotected lumen of the stomach to the accumulating acid [16-18] Pylorus ligation induced ulcers are shown by auto digestion of the gastric mucosa and breakdown of the gastric mucosal barrier which resulted as upper gastrointestinal damage including lesions, ulcers and life threatening perforation and haemorrhage. The pyloric ligation of the stomach causes accumulation of gastric acid which leads to development of ulceration in stomach. The agents who decrease gastric acid secretion and increase mucus secretion are effective in preventing the ulcers induced by this method. Like ranitidine, omeprazole acts as anti-ulcer agent by antisecretory mechanism via inhibition of gastric secretion and pepsin activity [12]. In the present study, ESIL prevents the ulcer may be by antisecretory and cytoprotective property.

Ethanol is responsible for disturbances in gastric secretion, damage to the mucosa, alterations in the permeability, gastric mucus depletion and free radical production. The generation of free radicals was produced by continuous release of superoxide

anion and hydroperoxy free radicals during metabolism of ethanol [19]. Ethanol induced gastric ulceration may be occurred due to stasis in gastric blood flow which contributes to the development of the haemorrhage and necrotic tissue injuries. Alcohol has ability to penetrate the gastric mucosa and causing the cellular damage which increases the permeability to sodium and water. In other hand, the accumulation of intracellular calcium causes the pathogenesis of gastric injury that leads to cell death and exfoliation of surface epithelium [16]. The present study observed that the ESIL significantly reduced ethanol induced ulcer by cytoprotective action via antioxidant effect. The ESIL extract showed cytoprotection against the ethanol and cysteamine induced ulceration by reducing the gastric acid secretion. Similarly, the cytoprotective effect of ESIL against cysteamine induced duodenal ulcers may be due to strengthening of duodenal mucosa or by increase of duodenal alkaline secretion or luminal prostaglandin level [12].

The results of this study found that ESIL established a cytoprotective action against ethanol induced cellular damage in the gastric mucosa of rats. Cytoprotection of anti-ulcer drugs has been recognised due to the generation of prostaglandins [20]. It has also been observed that ESIL significantly and dose dependently reduced the extent of gastric ulceration in pylorus ligated rats without affecting the gastric secretion or pepsin activity. The defence potential of mucus perimeter of gastric mucosa depends upon a delicate balance between the processes affecting the synthesis and secretion of mucin constituents. ESIL prevented the mucosal lesions induced by alcohol [21]. The modern approach towards a potent antiulcer agent involves a delicate balance of controlling the synthesis, secretion and metabolism of proteins, glycoproteins and lipids, so as to strengthen the mucosal integrity [22].

Several scientific studies revealed that the phytoconstituents like flavonoids, tannins, terpenoids and saponin were responsible for gastro protective agents [23]. Tannins possess as an antiulcer agent by its astringency property and vasoconstriction effects. Due to precipitation of micro proteins on the ulcer site, a protective layer was formed which hinders gut secretions and protects the mucosa from toxins and other irritants. Previous studies have recommended that these above active compounds had ability to stimulate mucus, bicarbonate and prostaglandin secretion and neutralize with the deteriorating effects of reactive oxidants in gastrointestinal lumen [24]. Therefore, ESIL possess antiulcer activity, may be due to presence of tanins, flavonoids and terpenoids.

CONCLUSION

The present study concluded that the antiulcer activity of ESIL may be attributed to antisecretory, cytoprotective and antioxidant properties. The bioactivity-guided phytochemical screening of ESIL revealed the presence of flavonoids, tannins and triterpenoids, which may be responsible for the anti-ulcer effect and can be further fractionated and investigated for their role and utility in any of the anti-ulcer mechanisms.

ACKNOWLEDGEMENTS

The authors would like to thank Dr. P. Venkateswara Rao, Principal and Head, Department of Pharmaceutical Chemistry, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India for providing laboratory facility and encouraging our work.

REFERENCES

- [1] Bharathi DP, Jegad E, Kavimani S. Antiulcer activity of aqueous extract of fruits of *Momordica cymbalaria* Hook f. in Wistar rats. *Pharmacognosy Res.* 2010;2(1):58-61.
- [2] Panneerselvam S, Arumugam G. A biochemical study on the gastroprotective effect of hydroalcoholic extract of *Andrographis paniculata* in rats. *Indian J Pharmacol.* 2011;43:402-08.
- [3] Arumugam S, Selvaraj SV, Velayutham S, Natesan SK, Palaniswamy K. Evaluation of anti-ulcer activity of *Samanea saman* (Jacq) merr bark on ethanol and stress induced gastric lesions in albino rats. *Indian J Pharmacol.* 2011;43:586-90.

- [4] Bandyopadhyay U, Biswas K, Chatterjee R, Bandyopadhyay D, Chattopadhyay I, Ganguly CK, et al. Gastroprotective effect of Neem (*Azadiracta indica*) bark extract possible involvement of H+K+ATPase inhibition and scavenging of hydroxyl radical. *Life Sci.* 2002;71:2845-65.
- [5] Anusha R. Herbal products in oral hygiene maintenance –a review. *IOSR Journal of Pharmacy.* 2015;5(1):48-51.
- [6] Nadkarni KM. Popular Prakashan, Mumbai, India, Indian Materia Medica. 1976;1:38-39.
- [7] Iyer D, Patil UK. Evaluation of antihyperlipidemic and antitumor activities of isolated coumarins from *Salvadora indica*. *Pharm Biol.* 2014 52(1):78-85.
- [8] Vimala G Gricilda SF. A review on antiulcer activity of few indian medicinal plants. *International Journal of Microbiology.* 2014;3(2):1-14.
- [9] David R. Katerere DL. Ethnoveterinary botanical medicine: herbal medicines for animal health, crc press, taylor and francis group, London New York. 2010; pp 293.
- [10] Khandelwal KR. Practical Pharmacognosy Techniques and Experiments. 2nd ed. Pune, India: Nirali Prakashan; 2000. Pp. 149-56.
- [11] OECD Test Guideline 425. Guidelines for Testing of Chemicals. Guidelines 425, Acute Oral Toxicity-Up-and-Down Procedure; 2001.
- [12] Deshpande SS, Shah GB, Parmar NS. Antiulcer activity of *Tephrosia purpurea* in rats. *Indian Journal of Pharmacology.* 2003;35:168-72.
- [13] Kulkarni SK. Hand book of experimental pharmacology, Vallabh Prakashan, New Delhi, 1999, Pp. 148-50.
- [14] Nie SN, Qian XM, Wu XH, Yang SY, Tang WJ, Xu BH, et al. Role of TFF in healing of stress induced gastric lesions. *World J Gastroenterol.* 2003;9:1772-76.
- [15] Robert A. Cytoprotection by prostaglandins. *Gastroenterol.* 1979;77:761-67.
- [16] Dwivedi V, Chander BS, Yadav NH. Evaluation of anti-ulcer activity of *Clitoria ternatea* Leaves (Linn) extract in Wistar rats. *Indian Journal of Research in Pharmacy and Biotechnology.* 2014;3:1225-29.
- [17] Raju D. Evaluation of Anti-ulcer activity of methanolic extract of *Terminalia chebula* fruits in experimental rats. *J Pharm Sci & Res.* 2009;3:101 07.
- [18] Salaj K, Mohammed A, Sunil SD, Satya P. Antiulcer activity of cod liver oil in rats, *Indian J Pharmacol.* 2008;40(5):209-14.
- [19] Jude EO, Paul A. Antiulcer and Anticonvulsant Activity of *Croton Zambesicus*. *J Pharm Sci.* 2009;22:384-90.
- [20] Devang JP, Indermeet SA. Anti-ulcer potential of *Oxystelma esculentum*. *International Journal of Green Pharmacy.* 2011;65-8.
- [21] Robert A, Nezmin JE, Lancaster C, Hanchar AJ. Cytoprotection by prostaglandins in rats. Prevention of gastric necrosis produced by HCl, NaOH, Hypertonic NaCl and thermal injury. *Gastroenterology.* 1979;76:439-43.
- [22] Brown GG. An introduction to Histotechnology, 1st edition. Appleton century Crofts, New York. 1978; Pp. 293-308.
- [23] Borelli F, Izzo AA. The plant kingdom as a source of anti-ulcer remedies. *Phytother Res.* 2000;14:581-91.
- [24] Sakat SS, Juvekar RA. Antiulcer activity of methanol extract of *erythrina indica* lam. leaves in experimental animals. *Pharmacognosy Research.* 2009;1: 396-401.

PARTICULARS OF CONTRIBUTORS:

1. Assistant Professor, Department of Pharmaceutical Chemistry, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.
2. Assistant Professor, Department of Pharmacology, MKCG Medical College & Hospital, Berhampur, Odisha, India.
3. Assistant Professor, Department of Pharmacology, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.
4. Assistant Professor, Department of Pharmaceutical Analysis, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.
5. Assistant Professor, Department of Pharmaceutics, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.
6. Assistant Professor, Department of Pharmacology, Sri Sivani College of Pharmacy, Srikakulam, Andhra Pradesh, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Himanshu Bhusan Sahoo,
Assistant Professor, Department of Pharmacology, MKCG Medical College & Hospital, Berhampur-760004, Odisha, India.
E-mail: bhusan.himanshu@yahoo.co.in

Date of Submission: **Mar 20, 2016**Date of Peer Review: **May 06, 2016**Date of Acceptance: **Jul 21, 2016**Date of Publishing: **Sep 01, 2016****FINANCIAL OR OTHER COMPETING INTERESTS:** None.